

PRESCRIBING INFORMATION

INCLUDING PATIENT MEDICATION INFORMATION

Pr Ampicillin Sodium for Injection BP

Powder for solution

250 mg, 500 mg, 1 g and 2 g per vial

Ampicillin (as ampicillin sodium)

For intramuscular or Intravenous Use

Sterile

Antibiotic

Auro Pharma Inc.
3700 Steeles Avenue West, Suite # 402
Woodbridge, Ontario, L4L 8K8
CANADA

Date of Revision:
September 16, 2019

Submission Control No: 231013

PRESCRIBING INFORMATION

Pr Ampicillin Sodium for Injection BP

Powder for solution

Antibiotic

Actions and Clinical Pharmacology

Ampicillin has a broad spectrum of bactericidal activity against many gram-positive and gram-negative aerobic and anaerobic bacteria. It acts through the inhibition of cell wall mucopeptide biosynthesis during the stage of active multiplication.

Indications and Clinical Use

The treatment of infections due to susceptible gram negative organisms (including strains of shigellae, *S. typhosa* and other salmonellae, *E. coli*, *H. influenzae*, and *P. mirabilis*) and susceptible gram positive organisms (including streptococci, pneumococci, and non-beta-lactamase (penicillinase) producing staphylococci).

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Ampicillin Sodium for Injection BP and other antibacterial drugs, Ampicillin Sodium for Injection BP should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

Contraindications

A history of allergic reactions to penicillin or cephalosporins.

Warnings

Before therapy, inquiry as to past penicillin or other allergies is essential as reactions occur more frequently in hypersensitive persons. During therapy, if allergic or anaphylactic reactions occur, discontinue treatment and initiate usual measures, i.e. antihistamines, pressor amines or corticosteroids. During long-term therapy, renal, hepatic, and hematopoietic functions should be checked periodically. Candidiasis and other superinfections may occur, especially in debilitated and malnourished patients, or those with low resistance to infection due to corticosteroids, immunosuppressors or irradiation.

The passage of any penicillin from blood into brain is facilitated by inflamed meninges and during cardiopulmonary bypass. In the presence of such factors and particularly in the presence of renal failure when high serum concentrations can be attained, central nervous system adverse effects including myoclonus, convulsive seizures and depressed consciousness can be expected. Although this complication has not been reported with ampicillin, it should be anticipated.

Precautions

General: A high percentage of patients with infectious mononucleosis or lymphatic leukemia who receive ampicillin develop a skin rash, and the drug should not be administered to such patients. In most cases, the rash is maculopapular, pruritic, and generalized. Prolonged use of antibiotics may promote overgrowth of nonsusceptible organisms. Should superinfections occur, appropriate measures should be taken.

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing Ampicillin Sodium for Injection BP in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of resistant organisms.

Drug Interactions

The concurrent administration of allopurinol and ampicillin increases substantially the incidence of rashes in patients receiving both drugs as compared to patients receiving ampicillin alone. It is not known whether this potentiation of ampicillin rashes is due to allopurinol or to hyperuricemia present in these patients. Ampicillin and aminoglycosides should not be reconstituted together due to the in vitro inactivation of the aminoglycosides by the ampicillin.

Drug/Laboratory Test Interactions: Following administration of ampicillin to pregnant women, a transient decrease in plasma concentration of total conjugated estriol, estriol glucuronide, conjugated estrone and estradiol has been noted.

With high urine concentrations of ampicillin, false-positive urinary glucose reactions may occur if copper reduction methods are used. Therefore, it is recommended that glucose tests based on enzymatic glucose oxidase reactions be employed.

Pregnancy: Animal studies with ampicillin have shown no teratogenic effects. There are, however, no adequate and well-controlled studies in pregnant women. Because animal reproduction studies are not always predictive of human response, this drug should be used during pregnancy only if clearly needed.

Lactation: Ampicillin is excreted in trace amounts in human milk. Therefore, caution should be exercised when ampicillin is administered to a nursing mother.

Use in Elderly: There are no known specific precautions for the use of ampicillin in the elderly.

Adverse Reactions

Gastrointestinal Disturbances: glossitis, stomatitis, black “hairy” tongue, nausea, vomiting, diarrhea, enterocolitis and pseudomembranous colitis. (These reactions are usually associated with oral administration.)

Hypersensitivity Reactions: Erythematous maculopapular rashes have been reported fairly frequently; urticaria, erythema multiforme, and a few cases of exfoliative dermatitis have been observed. Anaphylaxis is the most serious reaction usually associated with parenteral administration.

Note: Urticaria, other skin rashes, and serum sickness like reactions may be controlled with antihistamines, and if necessary, systemic corticosteroids. Serious anaphylactic reactions require the immediate use of epinephrine, oxygen and i.v. corticosteroids. In cases of infectious mononucleosis, where ampicillin has been administered, an extremely high incidence of generalized rash has been reported.

Renal: Interstitial nephritis has been reported.

Ototoxicity: Ampicillin may be ototoxic when given i.v. in very high doses.

Hepatic: A mild transitory elevation of serum glutamic oxaloacetic transaminase (SGOT) in individuals receiving large (2 to 4 times recommended dose) and often repeated i.m. injections.

Evidence indicates that serum glutamic oxaloacetic transaminase (SGOT) is released at the site of i.m. injection of sodium ampicillin and that the presence of the enzyme in the blood does not necessarily indicate liver involvement.

Hematologic Disturbances: Anemia, thrombocytopenia, thrombocytopenic purpura, hemorrhagic diathesis, eosinophilia, leukopenia and agranulocytosis have been reported rarely in association with ampicillin therapy. These reactions are usually reversible on discontinuation of the drug and are believed to be hypersensitivity phenomena.

Symptoms and Treatment of Overdosage

The treatment of overdosage would likely be needed only in patients with severely impaired renal function. In case of overdosage, discontinue medication, treat symptomatically and institute supportive measures as required. In patients with renal function impairment, ampicillin class antibiotics can be removed by hemodialysis but not by peritoneal dialysis.

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

Dosage and Administration

Dosage:

Infections of the ear, nose, throat and lower respiratory tract: **Adults:** 250 to 500 mg every 6 hours. **Children:** 25 to 50 mg/kg/day in equally divided doses at 6-hour intervals. Infections of gastrointestinal tract and of the genitourinary tract: **Adults:** 500 mg every 6 hours. **Children:** 50 mg/kg/day in equally divided doses at 6-hour intervals.

Larger doses may be required for stubborn or severe infections. The children's dosages are intended for individuals whose weights will not result in calculated dosage greater than that recommended for adults.

In the treatment of chronic urinary tract and intestinal tract infections, frequent bacteriological and clinical appraisal is necessary. Smaller doses than those recommended above should not be used; higher doses may be needed at times. In stubborn infections, therapy may be required for several weeks. It may be necessary to continue clinical and/or bacteriological follow-up for several months after cessation of therapy.

Treatment should be continued for a minimum of 48 to 72 hours beyond the time that the patient becomes asymptomatic or evidence of bacterial eradication has been obtained. A minimum of 10 days' treatment is recommended for any infection caused by Group A beta-hemolytic streptococci. In gonorrhea therapy, serologic tests for syphilis should be performed initially and monthly for 3 months.

Administration:

IM/Direct IV Use: Entire contents should be withdrawn and the dose injected over a period of 3-5 minutes.

IV Infusion: Entire contents should be withdrawn and the dose injected over a period of at least 10-15 minutes.

CAUTION: More rapid administration may result in convulsive seizures. The solution must be used immediately after reconstitution.

Pharmaceutical Information


Drug Substance

Proper name: Ampicillin Sodium

Chemical name: Sodium (2S,5R,6R)-6-[[[(2R)-2-amino-2-phenylacetyl]amino]-3,3-dimethyl-7-oxo-4-thia-1-azabicyclo[3.2.0]heptane-2-carboxylate.

Molecular formula and molecular mass: C₁₆H₁₈ N₃NaO₄S, 371.4 g/mol

Structural Formula:



Physicochemical properties: A White or almost white powder hygroscopic.

Stability & Storage Recommendations

Store the dry powder at room temperature (15°C to 25°C).

Reconstituted Solutions: Use sterile water for injection as the only diluent. Reconstituted solutions should be used immediately after reconstitution.

Reconstitution

I.M. Use: Using sterile water for injection, reconstitute as follows:

Vial Size (mg)	Volume of Diluent Added (mL)	Withdrawable Volume (mL)	Nominal Concentration (mg/mL)
250	1.9	2	125
500	1.8	2	250
1000	3.5	4	250

Withdraw the entire contents and use immediately after reconstitution.

Direct I.V. Use: Use sterile water for injection, reconstitute as follows:

Vial Size (mg)	Volume of Diluent Added (mL)	Withdrawable Volume (mL)	Nominal Concentration (mg/mL)
250	1.9	2	125
500	1.8	2	250

For direct intravenous administration, the product should be diluted to a concentration of 50 mg/mL with Sterile Water for Injection and administered by slow injection (three to four minutes).

Withdraw the entire contents and use immediately after reconstitution.

I.V. Infusion: Use sterile water for injection for initial dilution of the 1 g and 2 g vials, and reconstitute as follows:

Vial Size (mg)	Volume of Diluent Added (mL)	Withdrawable Volume (mL)	Nominal Concentration (mg/mL)
1000	3.5	4	250
2000	6.8	8	250

Withdraw the entire contents and use immediately after reconstitution.

Parenteral Products: Stability studies on ampicillin sodium, at concentrations of 2 mg/mL and 30 mg/mL in various i.v. solutions, indicate the drug will lose less than 10% activity at room temperature (22°C) for the time periods stated when the drug is added to the following infusion fluids:

Isotonic sodium chloride: (30 mg/mL) 8 hours
 5% dextrose in water: (2 mg/mL) 0 hours
 5% dextrose in 0.4% sodium chloride solution: (2 mg/mL) 0 hours

Availability of Dosage Forms:

Dosage form	Powder for Injection			
Strength	250 mg	500 mg	1 g	2 g
Description	White to off-white powder filled in 10 ml clear glass vials, stoppered with grey bromo butyl rubber stoppers and sealed with aluminium seals having taxim red coloured polypropylene disc.	White to off-white powder filled in 10 ml clear glass vials, stoppered with grey bromo butyl rubber stoppers and sealed with aluminium seals having violet coloured polypropylene disc.	White to off - white powder filled in 15 mL clear glass vials, stoppered with grey bromo butyl rubber stoppers and sealed with aluminium seals having grey coloured polypropylene disc.	White to off-white powder filled in 20 ml clear glass vials, stoppered with grey bromo butyl rubber stoppers and sealed with aluminium seals having yellow coloured polypropylene disc.
Composition	Each vial contains 282.5 mg of Ampicillin Sodium equivalent to 250 mg of Ampicillin. Non-medicinal Ingredients: water for injection	Each vial contains 564.9 mg of Ampicillin Sodium equivalent to 500 mg of Ampicillin. Non-medicinal Ingredients: water for injection	Each vial contains 1.13 g of Ampicillin Sodium equivalent to 1 g of Ampicillin. Non-medicinal Ingredients: water for injection	Each vial contains 2.26 g of Ampicillin Sodium equivalent to 2 g of Ampicillin. Non-medicinal Ingredients: water for injection
Packaging	10 mL clear glass vial. The available pack sizes are 10 x 10 Vials each.	10 mL clear glass vial. The available pack sizes are 10 x 10 Vials each.	15 mL clear glass vial. The available pack sizes are 10 x 10 Vials each.	20 mL clear glass vial. The available pack sizes are 10 x 10 Vials each.

REFERENCES:

- Prescribing information - ^{Pr} AMPICILLIN SODIUM FOR INJECTION USP. Teva Canada Limited., Control no. 216284 Date of Revision: December 17, 2018.

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

PATIENT MEDICATION INFORMATION

Pr Ampicillin Sodium for Injection BP

Powder for solution

Read this carefully before you start taking Ampicillin Sodium for Injection BP and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about Ampicillin Sodium for Injection BP.

What is Ampicillin Sodium for Injection BP used for:

Ampicillin Sodium for Injection BP is used to treat certain infections of the:

- Ear
- Nose
- Throat
- Lower Respiratory tract
- Antibacterial drugs like Ampicillin Sodium for Injection BP treat only bacterial infections. They do not treat viral infections.

How does Ampicillin Sodium for Injection BP work?

Ampicillin Sodium for Injection BP is an antibiotic that works by:

- Stopping the growth of bacteria.
- Killing bacteria.

What are the ingredients in Ampicillin Sodium for Injection BP?

Medicinal ingredients: Ampicillin Sodium

Ampicillin Sodium for Injection BP comes in the following dosage forms: Ampicillin Sodium for Injection BP is supplied as a dry powder in vials containing: 250 mg, 500 mg, 1 g or 2 g of ampicillin base as the sodium salt.

Do not use Ampicillin Sodium for Injection BP if:

- You have had an allergic reaction to Ampicillin Sodium for Injection BP or other medicines such as cephalosporins or penicillins.

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take Ampicillin Sodium for Injection BP. Talk about any health conditions or problems you may have, including if you:

- Have severe kidney disease with or without significant liver disease
- Are pregnant or planning to become pregnant
- Are breast feeding or planning to breastfeed.

Other warnings that you should know:

Ampicillin Sodium for Injection BP may affect certain urine test results. Remind your healthcare professional that you are taking Ampicillin Sodium for Injection BP if a urine test is ordered.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with Ampicillin Sodium for Injection BP:

- Allopurinol, used to treat gout or kidney stones

How to take Ampicillin Sodium for Injection BP:

- Although you may feel better early in treatment, Ampicillin Sodium for Injection BP should be used exactly as directed.
- Misuse or overuse of Ampicillin Sodium for Injection BP could lead to the growth of bacteria that will not be killed by Ampicillin Sodium for Injection BP (resistance). This means that Ampicillin Sodium for Injection BP may not work for you in the future.
- Do not share your medicine.

Usual Dose:

Adults: Your doctor will decide your dose based on your infection. The usual dose is 250 mg – 500 mg every 6 hours.

Children: Your doctor will decide your child’s dose based on your child’s weight and their infection.

Overdose:

If you think you have taken too much Ampicillin Sodium for Injection BP, contact your healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

What are possible side effects from using Ampicillin Sodium for Injection BP?

These are not all the possible side effects you may feel when taking Ampicillin Sodium for Injection BP. If you experience any side effects not listed here, contact your healthcare professional.

- Upper stomach pain
- Flatulence

Other side effects may occur that usually do not need medical attention. These side effects may go away during treatment as your body adjusts to the medicine. However, check with your doctor for any side effect that seems unusual or that is especially bothersome.

Serious side effects and what to do about them:

Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
Rare	an allergic reaction (difficulty in breathing, closing of the throat, swelling of the lips, face or tongue; hives or a rash)			✓
	redness, or itching			✓
	severe nausea, vomiting, or diarrhea			✓

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada/adverse-reaction-reporting.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

Dry Powder: Store at room temperature (15°C to 25°C).

Reconstituted Solutions: Use sterile water for injection as the only diluent. Reconstituted solutions should be used immediately after reconstitution.

Keep out of reach and sight of children.

If you want more information about Ampicillin Sodium for Injection BP:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://health-products.canada.ca/dpd-bdpp/index-eng.jsp>); the manufacturer’s website www.auropharma.ca, or by calling 1-855-648-6681.

This leaflet was prepared by:

Auro Pharma Inc.

3700 Steeles Avenue West, Suite # 402

Woodbridge, Ontario, L4L 8K8,

Canada.

Date of Revision: September 16, 2019.