

## **PRODUCT MONOGRAPH**

<sup>Pr</sup>**Auro-Telmisartan HCTZ**

Telmisartan and Hydrochlorothiazide Tablets, USP

80 mg/12.5 mg and 80 mg/25 mg

Angiotensin II AT1 Receptor Blocker/Diuretic

**Auro Pharma Inc.**  
3700 Steeles Avenue West, Suite # 402  
Woodbridge, Ontario, L4L 8K8,  
Canada.

Date of Revision:  
April 18, 2017

**Submission Control No: 204683**

**Table of Contents**

**PART I: HEALTH PROFESSIONAL INFORMATION..... 3**

SUMMARY PRODUCT INFORMATION ..... 3

INDICATIONS AND CLINICAL USE..... 3

CONTRAINDICATIONS ..... 4

WARNINGS AND PRECAUTIONS..... 4

ADVERSE REACTIONS..... 10

DRUG INTERACTIONS ..... 20

DOSAGE AND ADMINISTRATION ..... 25

OVERDOSAGE ..... 26

ACTION AND CLINICAL PHARMACOLOGY ..... 27

STORAGE AND STABILITY..... 32

DOSAGE FORMS, COMPOSITION AND PACKAGING ..... 32

**PART II: SCIENTIFIC INFORMATION ..... 33**

PHARMACEUTICAL INFORMATION..... 33

CLINICAL TRIALS ..... 35

DETAILED PHARMACOLOGY ..... 43

TOXICOLOGY ..... 44

REFERENCES ..... 47

**PART III: CONSUMER INFORMATION..... 48**

<sup>Pr</sup>**Auro-Telmisartan HCTZ**

Telmisartan and Hydrochlorothiazide Tablets, USP

80 mg/12.5 mg and 80 mg/25 mg

**PART I: HEALTH PROFESSIONAL INFORMATION**

**SUMMARY PRODUCT INFORMATION**

<b>Route of Administration</b>	<b>Dosage Form / Strength</b>	<b>All Nonmedicinal Ingredients</b>
oral	tablet 80/12.5 mg and 80/25 mg	Meglumine, povidone, sodium hydroxide pellets, colloidal silicon dioxide, mannitol, sodium stearyl fumarate, lactose monohydrate, hydroxypropyl cellulose, yellow ferric oxide (80 mg/25 mg) & red ferric oxide (80 mg/12.5 mg).

**INDICATIONS AND CLINICAL USE**

Auro-Telmisartan HCTZ (80 mg telmisartan/12.5 mg hydrochlorothiazide) is indicated for:

- treatment of mild to moderate essential hypertension in patients in whom combination therapy is considered appropriate.

Auro-Telmisartan HCTZ (80 mg telmisartan/25 mg hydrochlorothiazide) is indicated for:

- treatment of patients whose blood pressure is not adequately controlled by Auro-Telmisartan HCTZ 80 mg/ 12.5 mg or patients who have been previously stabilised on telmisartan and hydrochlorothiazide given separately.

These fixed-dose combinations are not indicated as initial therapy (see DOSAGE AND ADMINISTRATION).

***Geriatrics (> 65 years of age):***

No dosage adjustment is necessary. It should be recognized, however, that greater sensitivity in some older individuals cannot be ruled out.

***Pediatrics (< 18 years of age):***

Safety and efficacy of telmisartan and hydrochlorothiazide tablets have not been established in children and in adolescents up to 18 years.

## CONTRAINDICATIONS

Auro-Telmisartan HCTZ (telmisartan and hydrochlorothiazide) is contraindicated in:

- Concomitant use of angiotensin receptor antagonists (ARBs) –including the telmisartan component of Auro-Telmisartan HCTZ- with aliskiren-containing drugs in patients with diabetes mellitus (type 1 or type 2) or moderate to severe renal impairment (GFR < 60 ml/min/1.73m<sup>2</sup>) is contraindicated (see WARNINGS and PRECAUTIONS, **Cardiovascular**, **Dual Blockade of the Renin-Angiotensin System (RAS)** and **Renal**, and DRUG INTERACTIONS, **Dual Blockade of the Renin-Angiotensin System (RAS) with ARBs, ACEIs or aliskiren-containing drugs**).
- Patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container. For a complete listing, see the DOSAGE FORMS, COMPOSITION AND PACKAGING section of the product monograph
- Pregnant women (see WARNINGS AND PRECAUTIONS, Special Populations – Pregnant Women)
- Nursing women (see WARNINGS AND PRECAUTIONS, Special Populations – Nursing Women)
- Patients with anuria due to the presence of hydrochlorothiazide
- Patients with the rare hereditary condition of fructose intolerance
  - **Mannitol**: Auro-Telmisartan HCTZ tablets contain 477.18 mg of mannitol per maximum recommended daily dose.
- Patients with rare hereditary problems of galactose intolerance, e.g. galactosaemia, the Lapp Lactase deficiency or glucose-galactose malabsorption
  - **Lactose**: Auro-Telmisartan HCTZ tablets contain 29.60 mg of lactose monohydrate in the dose strength of 80 mg/12.5 mg and 80 mg/25 mg.

## WARNINGS AND PRECAUTIONS

### Serious Warnings and Precautions

**When used in pregnancy, angiotensin receptor (AT<sub>1</sub>) blockers (ARB) can cause injury or even death of the developing fetus. When pregnancy is detected, Auro-Telmisartan HCTZ should be discontinued as soon as possible (see WARNINGS AND PRECAUTIONS, Special Populations).**

### **General**

A case of rare but fatal angioedema occurred in a patient who had been medicated for about 6 months with telmisartan, one of the active components of telmisartan and hydrochlorothiazide tablets. The Autopsy Report described evidence of edema of the laryngeal mucosa, with terminal respiratory and circulatory failure. This is in the context of approximately 5.2 million patient-years exposure to telmisartan annually.

If laryngeal stridor or angioedema of the face, extremities, lips, tongue, or glottis occurs, Auro-Telmisartan HCTZ should be discontinued immediately, the patient treated appropriately in accordance with accepted medical care, and carefully observed until the swelling disappears. In

instances where swelling is confined to the face and lips, the condition generally resolves without treatment, although antihistamines may be useful in relieving symptoms. Where there is involvement of tongue, glottis, or larynx, likely to cause airway obstruction, appropriate therapy (including, but not limited to 0.3 to 0.5 ml of subcutaneous epinephrine solution 1:1000) should be administered promptly (see ADVERSE REACTIONS - Post Marketing Adverse Drug Reactions).

Patients with a known hypersensitivity (anaphylaxis) or angioedema to ARBs should not be treated with Auro-Telmisartan HCTZ (see ADVERSE REACTIONS, Clinical Trial Adverse Drug Reactions -All Clinical Trials, Immune System, Not known: angioedema and ADVERSE REACTIONS - Post Market Adverse Drug Reactions).

### **Carcinogenesis and Mutagenesis**

See TOXICOLOGY, Carcinogenicity and Mutagenicity.

### **Cardiovascular**

#### **Hypotension**

In patients who are volume-depleted by diuretic therapy, dietary salt restriction, dialysis, diarrhea or vomiting, symptomatic hypotension may occur after initiation of therapy with telmisartan. Such conditions, especially volume and/or sodium depletion, should be corrected prior to administration of telmisartan. In these patients, because of the potential fall in blood pressure, therapy with telmisartan should be initiated under close medical supervision. Similar considerations apply to patients with ischemic heart or cerebrovascular disease, in whom an excessive fall in blood pressure could result in myocardial infarction or cerebrovascular accident.

#### **Aortic and Mitral Valve Stenosis, Obstructive Hypertrophic Cardiomyopathy**

As with other vasodilators, special caution is indicated in patients suffering from aortic or mitral stenosis, or obstructive hypertrophic cardiomyopathy. These patients are at risk of decreased coronary perfusion resulting from a cardiac output that is limited by a fixed cardiac vascular obstruction.

#### **Dual blockade of the Renin-Angiotensin System (RAS)**

There is evidence that co- administration of angiotensin receptor antagonists (ARBs), such as the telmisartan component of telmisartan and hydrochlorothiazide tablets, or of angiotensin-converting-enzyme inhibitors (ACEIs) with aliskiren increases the risk of hypotension, syncope, stroke, hyperkalemia and deterioration of renal function, including renal failure, in patients with diabetes mellitus (type 1 or type 2) and/or moderate to severe renal impairment (GFR < 60 ml/min/1.73m<sup>2</sup>). Therefore, the use of Auro-Telmisartan HCTZ in combination with aliskiren-containing drugs is contraindicated in these patients (see CONTRAINDICATIONS).

Further, co-administration of ARBs, including the telmisartan component of Auro-Telmisartan HCTZ, with other agents blocking the RAS, such as ACEIs or aliskiren-containing drugs, is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, renal failure, and hyperkalemia.

## **Endocrine and Metabolism**

### **Electrolyte and Metabolic Imbalances**

#### *Telmisartan & Hydrochlorothiazide*

In controlled trials using telmisartan (80 mg) and hydrochlorothiazide (12.5 mg) in combination, there were no reports of hyperkalemia. Hypokalemia was reported in 1.4% of patients treated with the combination. No discontinuations due to hypokalemia occurred during treatment. The absence of significant changes in serum potassium levels may be due to the opposing mechanisms of action of telmisartan and hydrochlorothiazide on potassium excretion by the kidney.

The use of a dual renin-angiotensin-aldosterone system (RAAS) blockade may lead to increased occurrence of hyperkalemia when given as add-on therapy in patients with controlled blood pressure.

#### *Hydrochlorothiazide*

During thiazide diuretic therapy, periodic determinations of serum electrolytes to detect possible electrolyte imbalances should be performed at appropriate intervals. All patients receiving thiazide therapy should be observed for clinical signs of fluid or electrolyte imbalance, particularly hyponatremia, hypokalemia and hypochloremic alkalosis. Serum and urine electrolyte determinations are particularly important when the patient experiences excessive vomiting or receives parenteral fluids.

Hypokalemia may develop especially with brisk diuresis, when severe cirrhosis is present or after prolonged therapy. Interference with adequate oral electrolyte intake will also contribute to hypokalemia. Hypokalemia may induce cardiac arrhythmia and may also sensitize or exacerbate the response of the heart to the toxic effects of digitalis (e.g. increased ventricular irritability).

Although any chloride deficit is generally mild and usually does not require special treatment except under extraordinary circumstances (as in liver or renal disease), chloride replacement therapy may be required in the treatment of metabolic alkalosis.

Dilutional hyponatremia may occur in edematous patients in hot weather; appropriate therapy is water restriction rather than administration of salt, except in rare instances when the hyponatremia is life-threatening. In actual salt depletion, appropriate replacement is the therapy of choice.

Calcium excretion is decreased by thiazide diuretics which may cause intermittent and slight elevation of serum calcium in the absence of known disorders of calcium metabolism. Marked hypercalcemia may also be evidence of hyperparathyroidism. In the event of significant hypercalcemia, Auro-Telmisartan HCTZ should be discontinued followed by assessment of parathyroid function.

Thiazide diuretics have been shown to increase the urinary excretion of magnesium, which may result in hypomagnesemia.

Hyperuricemia may occur, and an acute attack of gout may be precipitated in certain patients

receiving thiazide therapy.

Insulin requirements in diabetic patients may be altered and latent diabetes mellitus may become manifest during thiazide diuretic therapy.

An increase in cholesterol and triglyceride levels has been associated with thiazide diuretic therapy.

Thiazide may decrease serum PBI levels without signs of thyroid disturbance.

### **Fertility**

No studies on fertility in humans have been performed (see Part II: TOXICOLOGY, Reproduction).

### **Hepatic**

#### **Hepatic Impairment**

As the predominant route of elimination of telmisartan is through biliary excretion, patients with cholestasis, biliary obstructive disorders or hepatic insufficiency have reduced clearance of telmisartan leading to increased systemic exposure. Auro-Telmisartan HCTZ should therefore be used with caution in these patients. Dosage reduction should be considered which would necessitate usage of the individual tablet formulations.

Auro-Telmisartan HCTZ is not recommended for patients with severe hepatic impairment. Thiazides should be used with caution in patients with impaired hepatic function or progressive liver disease, since minor alterations of fluid and electrolyte balance may precipitate hepatic coma (see DOSAGE AND ADMINISTRATION, Patients with Hepatic Impairment).

### **Neurologic**

#### **Effects on ability to drive and use machines**

No studies on the effect on the ability to drive and use machines have been performed. However, when driving vehicles or operating machinery it should be taken into account that dizziness or drowsiness may occasionally occur when taking antihypertensive therapy.

### **Immune**

#### **Hypersensitivity Reactions**

Hypersensitivity reactions to the hydrochlorothiazide component of Auro-Telmisartan HCTZ may occur in patients with or without a history of allergy or bronchial asthma.

#### **Systemic Lupus Erythematosus**

Thiazide diuretics have been reported to cause exacerbation or activation of systemic lupus erythematosus.

### **Renal**

#### **Renal Impairment**

In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney or severe congestive heart failure, dual blockade of the renin - angiotensin-aldosterone system

(e.g. concomitant use of an ARB with an ACE-inhibitor or the direct renin-inhibitor aliskiren) and treatment with agents that inhibit this system have been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk. Appropriate assessment of renal function should be conducted prior to use of Auro-Telmisartan HCTZ.

In studies of ACE inhibitors in patients with unilateral or bilateral renal artery stenosis, increases in serum creatinine or blood urea nitrogen were observed. Although there has been no long-term experience with telmisartan in this patient population, an effect similar to that observed with ACE inhibitors should be anticipated.

Due to the hydrochlorothiazide component, Auro-Telmisartan HCTZ is not recommended in patients with severe renal impairment (creatinine clearance  $\leq 30$  mL/min).

Thiazide diuretics should be used with caution in patients with renal impairment.

There is no experience regarding the administration of telmisartan and hydrochlorothiazide tablets in patients with a recent kidney transplant.

The use of ARBs – including the telmisartan component of Auro-Telmisartan HCTZ– or of ACEIs with aliskiren-containing drugs is contraindicated in patients with moderate to severe renal impairment (GFR  $< 60$  ml/min/1.73m<sup>2</sup>). (See CONTRAINDICATIONS and DRUG INTERACTIONS, Dual Blockade of the Renin-Angiotensin System (RAS) with ARBs, ACEIs, or aliskiren-containing drugs).

#### Azotemia

Azotemia may be precipitated or increased by the hydrochlorothiazide component of Auro-Telmisartan HCTZ. Cumulative effects of the drug may develop in patients with impaired renal function since the primary route of excretion is through the urine.

If increasing azotemia and oliguria occur during treatment of severe progressive renal disease, the diuretic should be discontinued.

#### Acute Myopia and Secondary Angle-Closure Glaucoma

Hydrochlorothiazide, a sulfonamide, has been associated with an idiosyncratic reaction resulting in acute transient myopia and acute angle-closure glaucoma. Symptoms include acute onset of decreased visual acuity or ocular pain and typically occur within hours to week of a drug initiation. Untreated acute-angle glaucoma can lead to permanent vision loss. The primary treatment is to discontinue hydrochlorothiazide as rapidly as possible. Prompt medical or surgical treatment may need to be considered if the intraocular pressure remains uncontrolled. Risk factors for developing acute angle closure glaucoma may include a history of sulfonamide or penicillin allergy.

#### Special Populations

##### **Pregnant Women:**

Drugs that act directly on the renin-angiotensin-aldosterone-system (RAAS) can cause fetal and neonatal morbidity and death when administered to pregnant women. When pregnancy is



detected, Auro-Telmisartan HCTZ should be discontinued as soon as possible.

The use of angiotensin receptor (AT<sub>1</sub>) blockers (ARBs) is not recommended during pregnancy and should not be initiated during pregnancy. Epidemiological evidence regarding the risk of teratogenicity following exposure to angiotensin converting enzyme inhibitors (another class of therapeutic products interfering with the RAAS) during the first trimester of pregnancy has not been conclusive; however a small increase in risk cannot be excluded. Given the current evidence available on the risk with ARB, similar risks may exist for this class of drugs. Patients planning pregnancy should be changed to alternative anti-hypertensive treatments which have an established safety profile for use in pregnancy. When pregnancy is diagnosed, treatment with angiotensin II antagonists should be stopped immediately, and, if appropriate, alternative therapy should be started.

The use of ARBs during the second and third trimesters is known to induce human fetotoxicity (decreased renal function, oligohydramnios, skull ossification retardation) and neonatal toxicity (renal failure, hypotension, hyperkalemia).

Infants with a history of in utero exposure to ARBs should be closely observed for hypotension, oliguria, and hyperkalemia. If oliguria occurs, attention should be directed toward support of blood pressure and renal perfusion. Exchange transfusion may be required as a means of reversing hypotension and/or substituting for disordered renal function; however, limited experience with those procedures has not been associated with significant clinical benefit.

There is limited experience with hydrochlorothiazide during pregnancy, especially during the first trimester. Hydrochlorothiazide crosses the placenta and appears in cord blood. Based on the pharmacological mechanism of action of hydrochlorothiazide its use during the second and third trimester may compromise feto-placental perfusion and may cause fetal and neonatal effects like icterus, disturbance of electrolyte balance, fetal or neonatal jaundice and thrombocytopenia.

Hydrochlorothiazide should not be used for gestational edema, gestational hypertension or preeclampsia due to the risk of decreased plasma volume and placental hypoperfusion, without a beneficial effect on the course of the disease. Hydrochlorothiazide should not be used for essential hypertension in pregnant women.

Diuretics do not prevent development of toxemia (preeclampsia) of pregnancy and there is no satisfactory evidence that they are useful in the treatment of toxemia.

There has been no clinical experience with telmisartan and hydrochlorothiazide tablets in pregnancy.

Preclinical studies with telmisartan do not indicate teratogenic effect, but have shown fetotoxicity.

**Nursing Women:** Telmisartan and hydrochlorothiazide tablets is contraindicated during lactation since it is not known whether telmisartan is excreted in human milk but significant levels have been found in the milk of lactating rats.

Animal studies have shown excretion of telmisartan in breast milk. Because many drugs are excreted in human milk and because of their potential for affecting the nursing infant adversely, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother. Thiazide diuretics are excreted in human milk at low levels.

**Diabetic Patients:**

In diabetic patients with undiagnosed coronary artery disease (CAD) on blood pressure lowering therapy, the risk of fatal myocardial infarction and unexpected cardiovascular death may be increased. In patients with diabetes mellitus, CAD may be asymptomatic and therefore undiagnosed. These patients should undergo appropriate diagnostic evaluation, e.g. exercise stress testing, to detect and to treat CAD accordingly before initiating blood pressure lowering treatment with Auro-Telmisartan HCTZ.

**Pediatrics (< 18 years of age):** Safety and effectiveness of telmisartan and hydrochlorothiazide tablets in pediatric patients have not been established.

**Geriatrics (> 65 years of age):** In clinical trials (n=1725) of patients treated with the combination of telmisartan and hydrochlorothiazide, 348 (20.2%) were 65 to 74 years of age and 78 (4.5%) were 75 years of age or older. No overall differences in the safety or efficacy profiles were observed in elderly patients compared with younger patients. It should be recognized however, that greater sensitivity of some older individuals cannot be ruled out.

**Monitoring and Laboratory Tests**

For specific monitoring and laboratory tests, see WARNINGS AND PRECAUTIONS (Cardiovascular, Endocrine and Metabolism, Hepatic and Renal) and DRUG INTERACTIONS sections.

**ADVERSE REACTIONS**

**Clinical Trial Adverse Drug Reactions**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.*

***Telmisartan and Hydrochlorothiazide Used in Combination***

The overall incidence and pattern of adverse events reported with telmisartan and hydrochlorothiazide tablets (80/25 mg) was comparable with telmisartan and hydrochlorothiazide tablets (80/12.5 mg). A dose-relationship of undesirable effects was not established and they showed no correlation with gender, age or race of the patients.

The combination of telmisartan and hydrochlorothiazide has been evaluated for safety in 1725

patients including 716 treated for over six months and 420 for over one year. In clinical trials with the individual components used in combination, no unexpected adverse events have been observed. Adverse experiences have been limited to those that have been previously reported with telmisartan and hydrochlorothiazide monotherapy. In general, treatment with the combination was well tolerated; most adverse experiences were mild and transient in nature and did not require discontinuation of therapy.

Adverse events at an incidence of  $\geq 1\%$  in patients treated with 80/12.5 mg telmisartan/hydrochlorothiazide combination, irrespective of their causal relationship, are presented in the following table. This table includes the results of two pivotal studies. One study, a factorial design, compared the use of various doses of telmisartan tablets and hydrochlorothiazide tablets in combination to telmisartan alone, hydrochlorothiazide alone and placebo. The other study compared the fixed dose combination 80/12.5 mg of telmisartan/hydrochlorothiazide to telmisartan 80 mg alone.

**Table 1: ADVERSE EVENTS OCCURRING IN  $\geq 1\%$  OF PATIENTS TREATED WITH 80/12.5 MG TELMISARTAN/HYDROCHLOROTHIAZIDE IN PIVOTAL CLINICAL TRIALS**

	Telmisartan/ HCTZ 80/12.5 mg (n=320) %	Telmisartan 80 mg (n=322) %	HCTZ 12.5 mg (n=75) %	Placebo (n=74) %
Total with any adverse event	39.1	41.3	46.7	41.9
Autonomic nervous system				
Sweating increased	1.3	0.3	0	0
Body as a whole				
Back Pain	1.6	2.5	1.3	0
Fatigue	2.8	2.2	4.0	1.4
Influenza-Like Symptoms	1.6	1.2	2.7	1.4
Pain	2.2	2.2	4.0	6.8
Central & peripheral nervous system				
Dizziness	6.9	3.7	2.7	1.4
Headache	2.5	4.0	13.3	16.2
Gastro-intestinal system				
Abdominal Pain	1.6	0.9	0	0
Diarrhoea	4.1	1.6	0	0
Nausea	1.6	0.9	1.3	0
Respiratory system				
Pharyngitis	1.6	0.3	0	0
Upper Respiratory Tract infection	2.5	3.7	9.3	6.8

HCTZ = hydrochlorothiazide

Note: Telmisartan 80 mg open label treatment is not included in the Telmisartan 80 mg column

Additional adverse reactions reported in clinical trials with telmisartan plus hydrochlorothiazide

are listed below according to system organ class:

**Autonomic Nervous System:** Impotence

**Body as a Whole:** Allergy, leg pain

**Central and Peripheral Nervous System:** Vertigo

**Gastro-intestinal System:** Dyspepsia, gastritis, gastro-intestinal disorder

**Metabolic and Nutritional System:** Hypokalaemia, loss of diabetic control, hyperuricaemia

**Musculo-Skeletal System:** Myalgia, arthralgia, arthrosis

**Nervous System Disorders:** sleep disturbances

**Psychiatric System:** Anxiety

**Respiratory System:** Bronchitis, sinusitis

**Skin and Appendages System:** Eczema, skin disorder

**Urinary System:** Urinary tract infection

In controlled trials with 1017 patients, 0.3% of patients treated with telmisartan (80 mg) and hydrochlorothiazide (12.5 mg) used in combination discontinued due to hypotension.

Adverse events occurred at approximately the same rates in men and women, older and younger patients and black and non-black patients.

#### **Abnormal Hematologic and Clinical Chemistry Findings**

In controlled trials, clinically relevant changes in standard laboratory test parameters were rarely associated with administration of telmisartan and hydrochlorothiazide in combination.

**Table 2: Laboratory Parameter Results in Patients Treated with Telmisartan and Hydrochlorothiazide in Combination**

<b>Laboratory Parameter</b>	<b>% of Patients Treated with Telmisartan/ Hydrochlorothiazide</b>	<b>Clinical Comment</b>
Increases in Blood Urea Nitrogen (BUN) ( $\geq 11.2$ mg/dL)	2.8%	No patient discontinued treatment due to an increase in BUN.
Increases in Serum Creatinine ( $\geq 0.5$ mg/dL)	1.4%	No patient discontinued treatment due to an increase in creatinine.

Laboratory Parameter	% of Patients Treated with Telmisartan/ Hydrochlorothiazide	Clinical Comment
Decreases in Hemoglobin ( $\geq 2$ g/dL)	1.2%	Changes in hemoglobin were not considered clinically significant and there were no discontinuations due to anemia.
Decreases in Hematocrit ( $\geq 9\%$ )	0.6%	Changes in hematocrit were not considered clinically significant and there were no discontinuations due to anemia.

**Liver function Tests:** Occasional elevations of liver enzymes and/or serum bilirubin have occurred. No telmisartan/hydrochlorothiazide treated patients discontinued therapy due to abnormal liver function.

**Serum Electrolytes:** see WARNINGS AND PRECAUTIONS.

### ***Telmisartan***

Additional side effects were reported in clinical trials with telmisartan in the indication hypertension or in patients 50 years or older at high risk of cardiovascular events.

Telmisartan has been evaluated for safety in 27 clinical trials involving 7968 patients. Of these 7968 patients, 5788 patients were treated with telmisartan monotherapy including 1058 patients treated for  $\geq 1$  year and 1395 patients treated in placebo-controlled trials.

The following potentially serious adverse events have been reported rarely with telmisartan in controlled clinical trials: syncope and hypotension. In placebo-controlled trials, no serious adverse event was reported with a frequency  $\geq 0.1\%$  in telmisartan-treated patients.

### **All Clinical Trials**

The adverse drug events listed below have been accumulated from 27 clinical trials including 5788 hypertensive patients treated with telmisartan. Adverse events have been ranked under headings of frequency using the following convention: very common ( $\geq 1/10$ ); common ( $\geq 1/100$ ,  $< 1/10$ ); uncommon ( $\geq 1/1000$ ,  $< 1/100$ ); rare ( $\geq 1/10000$ ,  $< 1/1000$ ); very rare ( $< 1/10000$ )

### **Body as a Whole, General:**

- Common: Chest pain, influenza-like symptoms, symptoms of infection (e.g. urinary tract infection including cystitis), fatigue, conjunctivitis.
- Uncommon: Hyperhidrosis, asthenia (weakness).

### **Blood and Lymphatic System:**

- Uncommon: Anemia.
- Rare: Thrombocytopenia.
- Unknown: eosinophilia.

### **Cardiovascular System:**

- Common: Edema, palpitation.

Uncommon: Bradycardia, orthostatic hypotension, hypotension.  
Rare: Tachycardia.

**Central and Peripheral Nervous System:**

Very Common: Headache.  
Common: Dizziness, insomnia.  
Uncommon: Vertigo.

**Eye Disorders:**

Rare: Visual disturbance.

**Gastro-Intestinal System:**

Common: Abdominal pain, diarrhoea, dyspepsia, nausea, constipation, gastritis.  
Uncommon: Dry mouth, flatulence, vomiting.  
Rare: Stomach discomfort.

**Hepato-biliary Disorders:**

Rare: Hepatic function abnormal/liver disorder\*.  
\*Most cases of hepatic function abnormal / liver disorder from post-marketing experience with telmisartan occurred in patients in Japan, who are more likely to experience these adverse reactions

**Immune System:**

Rare: Hypersensitivity, Exacerbation or activation of systemic lupus erythematosus\*  
\* based on post-marketing experience.  
Unknown: Anaphylactic reaction, angioedema.

**Infections and Infestations:**

Uncommon: Upper respiratory tract infections, urinary tract infections.  
Not known: Sepsis including fatal outcome.

**Investigations:**

Uncommon: Blood creatinine increased.  
Rare: Blood uric acid increased, hepatic enzymes increased, blood creatinine phosphokinase increased, haemoglobin decreased.

**Metabolism and Nutrition Disorders:**

Uncommon: Hyperkalemia.  
Rare: Hypoglycemia (in diabetic patients)

**Musculo-Skeletal System:**

Common: Arthralgia, muscle spasms (cramps in legs) or pain in extremity (leg pain), myalgia, arthritis, arthrosis.  
Uncommon: Tendon pain (tendonitis like symptoms), back pain.

**Nervous System:**

Uncommon: Syncope (faint).

**Psychiatric System:**

Common: Anxiety, nervousness.

Uncommon: Depression.

**Renal and Urinary System:**

Uncommon: Renal impairment including acute renal failure.

**Respiratory System:**

Common: Upper respiratory tract infections including pharyngitis and sinusitis, bronchitis, coughing, dyspnea, rhinitis.

**Skin and Appendages System:**

Common: Skin disorders like rash.

Uncommon: Pruritus.

Rare: Erythema, drug eruption, eczema, toxic skin eruption.

Unknown: Urticaria.

**Placebo-Controlled Trials**

The overall incidence of adverse events reported with telmisartan (41.4%) was usually comparable to placebo (43.9%) in controlled clinical trials.

Adverse events occurring in  $\geq 1\%$  of 1395 hypertensive patients treated with telmisartan *monotherapy* in placebo-controlled clinical trials, regardless of drug relationship, include the following:

**Table 3: Adverse Events Occurring in  $\geq 1\%$  of 1395 Hypertensive Patients Treated with Telmisartan Monotherapy**

Adverse Event, by System	Telmisartan Total N=1395 %	Placebo N=583 %
Body as a Whole		
Back Pain	2.7	0.9
Chest Pain	1.3	1.2
Fatigue	3.2	3.3
Influenza-Like Symptoms	1.7	1.5
Pain	3.5	4.3
Central & Peripheral Nervous System		
Dizziness	3.6	4.6
Headache	8.0	15.6
Somnolence	0.4	1.0
Gastrointestinal System		
Diarrhea	2.6	1.0

<b>Adverse Event, by System</b>	<b>Telmisartan Total N=1395 %</b>	<b>Placebo N=583 %</b>
Dyspepsia	1.6	1.2
Nausea	1.1	1.4
Vomiting	0.4	1.0
<b>Musculoskeletal System</b>		
Myalgia	1.1	0.7
<b>Respiratory System</b>		
Coughing	1.6	1.7
Pharyngitis	1.1	0.3
Sinusitis	2.2	1.9
Upper Respiratory Tract Infection	6.5	4.6
<b>Heart Rate and Rhythm Disorders</b>		
ECG abnormal specific	0.2	1.0
Palpitation	0.6	1.0
<b>Cardiovascular Disorders, General</b>		
Hypertension	1.0	1.7
Oedema peripheral	1.0	1.2

The incidence of adverse events was not dose-related and did not correlate with the gender, age, or race of patients.

**Less Common Clinical Trial Adverse Drug Reactions (<1%)**

In addition, the following adverse events, with no established causality, were reported at an incidence <1% in placebo-controlled clinical trials:

**Autonomic Nervous System Disorders:** sweating increased.

**Body as a Whole:** abdomen enlarged, allergy, cyst nos, fall, fever, leg pain, rigors, syncope.

**Cardiovascular Disorders, General:** hypotension, hypotension-postural, leg edema.

**Central & Peripheral Nervous System Disorder:** hypertonia, migraine-aggravated, muscle contraction-involuntary.

**Gastrointestinal System Disorders:** anorexia, appetite increased, flatulence, gastrointestinal disorder nos, gastroenteritis, gastroesophageal reflux, melena, mouth dry, abdominal pain.

**Heart Rate & Rhythm Disorders:** arrhythmia, tachycardia.

**Metabolic & Nutritional Disorders:** diabetes mellitus, hypokalaemia.



**Musculoskeletal System Disorders:** arthritis, arthritis aggravated, arthrosis, bursitis, fascitis plantar, tendon pain.

**Myo Endo Pericardial & Valve Disorders:** myocardial infarction.

**Psychiatric Disorders:** nervousness.

**Red Blood Cell Disorders:** anemia.

**Reproductive Disorders, Female:** vaginitis.

**Resistance Mechanism Disorders:** abscess, infection, bacterial, moniliasis genital, otitis media.

**Respiratory System Disorders:** bronchospasm, epistaxis, pneumonia, bronchitis.

**Skin & Appendage Disorders:** rash, skin dry.

**Urinary System Disorders:** dysuria, hematuria, micturition disorder, urinary tract infection (including cystitis).

**Vascular (Extracardiac) Disorders:** cerebrovascular disorder, purpura.

**Vision Disorders:** vision abnormal.

Angioedema has been reported rarely in patients treated with telmisartan.

**Abnormal Hematologic and Clinical Chemistry Findings**

In placebo-controlled clinical trials involving 1041 patients treated with telmisartan monotherapy, clinically relevant changes in standard laboratory test parameters were rarely associated with administration of telmisartan.

**Table 4: Laboratory Parameter Results in Placebo-Controlled Clinical Trials Involving 1041 Patients Treated With TELMISARTAN Monotherapy**

<b>Laboratory Parameter</b>	<b>% of Placebo Patients</b>	<b>% of Patients Treated with Telmisartan</b>	<b>Clinical Comment</b>
Increases in ALT > 3 times the upper limit of normal	1.7%	0.5%	No telmisartan-treated patients discontinued therapy due to abnormal hepatic function.
Increases in AST > 3 times the upper limit of normal	0.8%	0.1%	No telmisartan-treated patients discontinued therapy due to abnormal hepatic function.
Increases in Blood Urea Nitrogen (BUN) ≥ 11.2 mg/dl	0.3%	1.5%	These increases occurred primarily with telmisartan in combination with hydrochlorothiazide. One telmisartan treated patient discontinued therapy due to increases in

Laboratory Parameter	% of Placebo Patients	% of Patients Treated with Telmisartan	Clinical Comment
			blood urea nitrogen and creatinine.
Increases in Creatinine $\geq 0.5$ mg/dl	0.3%	0.6%	These increases occurred primarily with telmisartan in combination with hydrochlorothiazide. One telmisartan treated patient discontinued therapy due to increases in blood urea nitrogen and creatinine.
Increases in Serum Potassium ( $\geq 1.4$ mEq/L)	0.6%	0.3%	Clinically significant changes in potassium (that exceeded 3 mEq/L) were found in 0.6% of telmisartan treated patients, with 0.5% of these reported at baseline. The corresponding rates for placebo-treated patients were 0.6% and 0.8%.
Decreases in Serum Potassium ( $\geq 1.4$ mEq/L)	0.3%	0.1%	Clinically significant changes in potassium (that exceeded 3 mEq/L) were found in 0.6% of telmisartan-treated patients, with 0.5% of these reported at baseline. The corresponding rates for placebo-treated patients were 0.6% and 0.8%.
Increases in Serum Uric Acid $\geq 2.7$ mg/dl	0.0%	1.7%	Clinically significant hyperuricemia ( $>10$ mEq/L) was observed in 2.3% of patients with telmisartan, with 0.4% reported in patients at baseline. Increases in serum uric acid were primarily observed in patients who received telmisartan in combination with hydrochlorothiazide. No patient was discontinued from treatment due to hyperuricemia.

### Hemoglobin, Hemotocrit

Clinically significant changes in hemoglobin and hematocrit ( $<10$ g/dl and  $<30\%$ , respectively) were rarely observed with telmisartan treatment and did not differ from rates in placebo-treated patients. No patients discontinued therapy due to anemia.

### Cholesterol

In placebo-controlled trials, marked increases in serum cholesterol were reported in a total of 6 telmisartan-treated patients (0.4%) and no placebo patients. Two of these patients were followed over time, in both cases cholesterol values reverted to baseline levels.

Serum elevations in cholesterol were reported as adverse events in 11 of 3445 patients (0.3%) in all clinical trials. There were no reported cases of hypercholesterolemia in telmisartan-treated patients in placebo-controlled trials.

## **Other Clinical Trials**

**Gastrointestinal:** constipation

**Respiratory:** rhinitis, dyspnea

**Special senses:** conjunctivitis

## **Post-Market Adverse Drug Reactions**

### ***Telmisartan***

Since the introduction of telmisartan in the market, cases of anxiety, dizziness, vision troubled, vertigo, abdominal distension, abdominal pain, retching, hyperhidrosis, arthralgia, myalgia, muscle spasm, back pain, asthenia, pain in extremity, fatigue, chest pain, blood creatinine increased, erythema, pruritus, syncope/faint, insomnia, depression, stomach discomfort, vomiting, hypotension (including orthostatic hypotension), bradycardia, tachycardia, abnormal hepatic function/liver disorder, renal impairment including acute renal failure, hyperkalemia, dyspnoea, anaemia, eosinophilia, thrombocytopenia and weakness have been reported. The frequency of these effects is unknown. As with other angiotensin II antagonists, rare cases of angioedema (with fatal outcome), pruritus, rash and urticaria have been reported.

Cases of muscle pain, muscle weakness, myositis and rhabdomyolysis have been reported in patients receiving angiotensin II receptor blockers.

In addition, since the introduction of telmisartan in the market, cases with increased blood creatinine phosphokinase (CPK) have been reported.

### ***Hydrochlorothiazide***

Adverse experiences that have been reported with hydrochlorothiazide alone without regard to causality are listed below:

**Body as a Whole:** fever

**Gastro-Intestinal System:** pancreatitis, sialadenitis, gastric irritation, anorexia, nausea, vomiting, diarrhea, constipation;

**Hepatobiliary Disorders:** Jaundice (hepatocellular cholestatic jaundice).

**Blood and Lymphatic System:** aplastic anemia, agranulocytosis, leukopenia, hemolytic anemia, thrombocytopenia (sometimes with purpura), bone marrow depression;

**Hypersensitivity:** purpura

**Respiratory Disorders:** respiratory distress including pneumonia and pulmonary edema

**Musculoskeletal:** muscle spasm, weakness, cramps in legs

**Metabolism and Nutrition Disorders:** volume depletion, loss of appetite, hypomagnesemia, hypercalcemia, hypochloremic alkalosis.

**Nervous System Disorders:** Headache, light-headedness.

**Central and Peripheral Nervous System:** dizziness, vertigo, paraesthesia, restlessness, nervousness;

**Cardiovascular:** orthostatic hypotension;

**Heart rate and rhythm disorder:** cardiac arrhythmias;

**Renal:** renal failure, renal dysfunction, interstitial nephritis;

**Skin and Subcutaneous Tissue Disorders:** rash, urticaria, erythema multiforme including Stevens-Johnson syndrome, exfoliative dermatitis including toxic epidermal necrolysis, photosensitivity reactions, necrotizing angiitis (vasculitis, cutaneous vasculitis), anaphylactic reactions, cutaneous lupus erythematosus-like reactions, reactivation of cutaneous lupus erythematosus

**Eye Disorders:** transient blurred vision, xanthopsia, acute myopia, acute angle-closure glaucoma.

**Laboratory Findings:** Metabolic: hyperglycaemia, glucosuria, hyperuricaemia;

**Other:** electrolyte imbalances (including hyponatraemia and hypokalaemia), increases in triglycerides, hypercholesterolemia.

## **DRUG INTERACTIONS**

### **Overview**

#### ***Telmisartan***

**Cytochrome P450:** Telmisartan is not metabolized by the cytochrome P450 (CYP) isoenzymes; as such, it is not expected that a pharmacokinetic interaction of telmisartan with drugs which inhibit or induce CYP isozymes will occur.

#### ***Hydrochlorothiazide***

**Cytochrome P450:** Hydrochlorothiazide is not metabolized by humans; as such, no pharmacokinetic interaction with agents known to inhibit or induce CYP isozymes or other enzymes systems is expected.

## Drug-Drug Interactions

**Table 5 - Established or Potential Drug-Drug Interactions**

Telmisartan	Effect	Clinical comment
Diuretics	Patients on diuretics, and especially those in whom diuretic therapy was recently instituted, may occasionally experience an excessive reduction of blood pressure after initiation of therapy with telmisartan.	The possibility of symptomatic hypotension with the use of telmisartan can be minimized by discontinuing the diuretic prior to initiation of treatment and/or lowering the initial dose of telmisartan (see WARNINGS - Hypotension and DOSAGE AND ADMINISTRATION). No drug interaction of clinical significance has been identified with thiazide diuretics.
Agents increasing serum potassium		Since the telmisartan component of Auro-Telmisartan HCTZ reduces the production of aldosterone, potassium-sparing diuretics or potassium supplements should be given only for documented hypokalemia and with frequent monitoring of serum potassium. Potassium-containing salt substitutes should also be used with caution. Concomitant thiazide diuretic use may attenuate any effect that telmisartan may have on serum potassium.
Digoxin	When telmisartan was co-administered with digoxin, mean increases in digoxin peak plasma concentration (49%) and in trough concentration (20%) were observed.	It is recommended that digoxin levels be monitored with appropriate dose adjustments when initiating, adjusting or discontinuing Auro-Telmisartan HCTZ, to maintain appropriate plasma digoxin concentrations.
Dual Blockade of the Renin-Angiotensin System (RAS) with ARBs, ACEIs or aliskiren-containing drugs		Dual Blockade of the renin-angiotensin system with ARBs, ACEIs or aliskiren-containing drugs is contraindicated in patients with diabetes and/or renal impairment, and is generally not recommended in other patients, since such treatment has been associated with an increased incidence of severe hypotension, renal failure, and hyperkalemia. (See CONTRAINDICATIONS and WARNINGS AND PRECAUTIONS, <u>Dual Blockade of the Renin-Angiotensin System</u> ).
Lithium Salts	Reversible increases in serum lithium concentrations and toxicity have been reported during concomitant administration of lithium with angiotensin converting enzyme inhibitors. Rare cases have also been reported with angiotensin II receptor antagonists including telmisartan.	Serum lithium level monitoring is advisable during concomitant use.

<b>Telmisartan</b>	<b>Effect</b>	<b>Clinical comment</b>
Nonsteroidal Anti-Inflammatory Drugs (NSAIDs)	Combinations of angiotensin-II antagonists (telmisartan) and NSAIDs (including ASA and COX-2 inhibitors) might have an increased risk for acute renal failure and hyperkalemia. NSAIDs (including ASA and COX-2 inhibitors) and angiotensin-II receptor antagonists exert a synergistic effect on the decrease of glomerular filtration. In patients with pre-existing renal impairment, this may lead to acute renal failure.	Blood pressure and kidney function should be monitored more closely in this situation, as occasionally there can be a substantial increase in blood pressure. Monitoring of renal function at the beginning and during the course of the treatment should be recommended. Co-administration of telmisartan did not result in a clinically significant interaction with ibuprofen.
Ramipril	In one study, the co-administration of telmisartan and ramipril led to an increase of up to 2.5 fold in the AUC <sub>0-24</sub> and C <sub>max</sub> of ramipril and ramiprilat.	The clinical relevance of this observation is not known.
Warfarin	Telmisartan administered for 10 days slightly decreased the mean warfarin trough plasma concentration; this decrease did not result in a change in the International Normalized Ratio (INR).	
Acetaminophen, amlodipine, glibenclamide, hydrochlorothiazide, or ibuprofen		Co-administration of telmisartan did not result in a clinically significant pharmacokinetic interaction

**Table 6 - Established or Potential Drug-Drug Interactions**

<b>Hydrochlorothiazide</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Alcohol, barbiturates and narcotics	C	Potential of orthostatic hypotension may occur.	Avoid alcohol, barbiturates or narcotics, especially with initiation of therapy.
Amphotericin B	T	Amphotericin B increases the risk of hypokalemia induced by thiazide diuretics.	Monitor of serum potassium level.
Anti-diabetic drugs (e.g. oral hypoglycemic agents and insulin)	CT	Thiazide-induced hyperglycemia may compromise blood sugar control. Depletion of serum potassium augments glucose intolerance.	Monitor glycemic control, supplement potassium if necessary, to maintain appropriate serum potassium levels, and adjust diabetes medications as required.
Antihypertensive drugs	CT	Hydrochlorothiazide may potentiate the action of other antihypertensive drugs (e.g. guanethidine, methyl dopa, beta-blockers, vasodilators, calcium channel blockers, ACEI, ARB, and direct renin inhibitors).	

<b>Hydrochlorothiazide</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Antineoplastic drugs, including cyclophosphamide and methotrexate	C	Concomitant use of thiazide diuretics may reduce renal excretion of cytotoxic agents and enhance their myelosuppressive effects.	Hematological status should be closely monitored in patients receiving this combination. Dose adjustment of cytotoxic agents may be required.
Bile acid sequestrants, e.g. cholestyramine and colestipol resins	CT	Bile acid sequestrants bind thiazide diuretics in the gut and impair gastrointestinal absorption by 43-85%. Administration of thiazide 4 hours after a bile acid sequestrant reduced absorption of hydrochlorothiazide by 30-35%.	Give thiazide 2-4 hours before or 6 hours after the bile acid sequestrant. Maintain a consistent sequence of administration. Monitor blood pressure, and increase dose of thiazide, if necessary.
Calcium and vitamin D supplements	C	Thiazides decrease renal excretion of calcium and increase calcium release from bone.	Monitor serum calcium, especially with concomitant use of high doses of calcium supplements. Dose reduction or withdrawal of calcium and/or vitamin D supplements may be necessary.
Carbamazepine	C	Carbamazepine may cause clinically significant hyponatremia. Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Corticosteroids, and adrenocorticotropic hormone (ACTH)	T	Intensified electrolyte depletion, particularly hypokalemia may occur.	Monitor serum potassium, and adjust medications, as required.
Digoxin	CT	Thiazide-induced electrolyte disturbances, e.e. hypokalemia, hypomagnesemia, increase the risk of digoxin toxicity, which may lead to fatal arrhythmic events.	Concomitant administration of hydrochlorothiazide and digoxin requires caution. Monitor electrolytes and digoxin levels closely. Supplement potassium or adjust doses of digoxin or thiazide, as required.
Drugs that alter GI motility, i.e., anticholinergic agents, such as atropine and prokinetic agents, such as metoclopramide, domperidone	CT, T	Bioavailability of thiazide diuretics may be increased by anticholinergic agents due to a decrease in gastrointestinal motility and gastric emptying. Conversely, prokinetic drugs may decrease the bioavailability of thiazide diuretics.	Dose adjustment of thiazide may be required.
Gout medications (allopurinol, uricosurics, xanthine oxidase inhibitors)	T, RCS	Thiazide-induced hyperuricemia may compromise control of gout by allopurinol and probenecid. The co-administration of hydrochlorothiazide and allopurinol may increase the incidence of hypersensitivity reactions to allopurinol.	Dosage adjustment of gout medications may be required.
Lithium	CT	Thiazide diuretics reduce the renal clearance of lithium and add a high risk of lithium toxicity.	Concomitant use of thiazide diuretics with lithium is generally not recommended. If such use is deemed necessary, reduce lithium dose by 50% and monitor lithium levels closely.

<b>Hydrochlorothiazide</b>	<b>Ref.</b>	<b>Effect</b>	<b>Clinical comment</b>
Non-steroidal Anti-Inflammatory drugs (NSAIDs including ASA and COX-2 inhibitors)	CT	The co-administration of a non-steroidal anti-inflammatory agent can reduce the diuretic, natriuretic, and antihypertensive effects of loop, potassium-sparing and thiazide diuretics. The potential for acute renal insufficiency in patients who are dehydrated may be enhanced. NSAID-related retention of sodium and water antagonises the diuretic and antihypertensive effects of thiazides. NSAID-induced inhibition of renal prostaglandins leading to decreases of renal blood flow, along with thiazide-induced decreases in GFR may lead to acute renal failure. Patients with heart failure may be at particular risk.	Patients receiving NSAIDs and Auro-Telmisartan HCTZ should be adequately hydrated and be monitored for renal function at the beginning of the combined treatment. Monitoring of renal function at the beginning and during the course of the treatment is recommended as well as regular hydration of the patient. Therefore, when Auro-Telmisartan HCTZ and NSAIDs are used concomitantly, the patient should be observed closely to determine whether the desired effect of the diuretic is obtained. If combination use is necessary, also monitor serum potassium and blood pressure closely. Dose adjustments may be required.
Pressor amines (e.g. norepinephrine)		Decreased response to pressor amines may occur, but the effect is considered not sufficient to preclude their concurrent use.	
Selective serotonin reuptake inhibitors (SSRIs, e.g. citalopram, escitalopram, sertraline)	T, C	Concomitant use with thiazide diuretics may potentiate hyponatremia.	Monitor serum sodium levels. Use with caution.
Skeletal muscle relaxants of the curare family e.g. tubocurarine	C	Thiazide drugs may increase the responsiveness of some skeletal muscle relaxants, such as curare derivatives.	
Topiramate	CT	Additive hypokalemia. Possible thiazide-induced increase in topiramate serum concentrations.	Monitor serum potassium and topiramate levels. Use potassium supplements, or adjust topiramate dose as necessary.
$\beta$ -adrenergic receptor blocking agents propranolol, metoprolol, sotalol, or acebutolol			No significant pharmacokinetic interactions were noted when these agents were administered concomitantly, separately or in fixed combination.
Spironolactone, indomethacin, allopurinol and phenytoin.			No significant interactions have been noted.

Legend: C = Case Study; RCS = Retrospective Cohort Study; CT = Clinical Trial; T = Theoretical

### **Drug-Food Interactions**

When telmisartan is taken with food, the reduction in the area under the plasma concentration-time curve (AUC) of telmisartan varies from approximately 6% (40 mg) to approximately 19% (160 mg), and the reduction in  $C_{max}$  varies from approximately 26% (40 mg) to 56% (160 mg). However, three hours after administration, plasma concentrations are similar whether telmisartan



is taken with or without food.

### **Drug-Herb Interactions**

Interactions with herbal products have not been established.

### **Drug-Laboratory Interactions**

Interactions with laboratory tests have not been established.

### **Drug-Lifestyle Interactions**

Interactions with lifestyle have not been established.

## **DOSAGE AND ADMINISTRATION**

### **Dosing Considerations**

Auro-Telmisartan HCTZ may be substituted in patients who have been stabilized on the individual telmisartan 80 mg and hydrochlorothiazide 12.5 mg components as described below.

Auro-Telmisartan HCTZ may be administered with or without food, however it should be taken consistently with regard to food intake.

### **Recommended Dose and Dosage Adjustment**

Auro-Telmisartan HCTZ (telmisartan/hydrochlorothiazide) is not for initial therapy.

A patient whose blood pressure is not adequately controlled with telmisartan monotherapy 80 mg, may be switched to Auro-Telmisartan HCTZ, (telmisartan 80 mg/hydrochlorothiazide 12.5 mg) once daily.

A patient whose blood pressure is not adequately controlled with Auro-Telmisartan HCTZ (telmisartan 80 mg and hydrochlorothiazide 12.5 mg tablets), may be switched to Auro-Telmisartan HCTZ (telmisartan 80 mg and hydrochlorothiazide 25 mg tablets) once daily.

### **Telmisartan Monotherapy**

The recommended dose of telmisartan is 80 mg once daily. The antihypertensive effect is present within 2 weeks and maximal reduction is generally attained after four weeks. If additional blood pressure reduction is required, a thiazide diuretic may be added.

No initial dosing adjustment is necessary for elderly patients or for patients with renal impairment but greater sensitivity in some older individuals cannot be ruled out. Markedly reduced telmisartan plasma levels were observed in patients on hemodialysis.

### **Diuretic Treated Patients**

In patients receiving diuretics, telmisartan therapy should be initiated with caution, since these patients may be volume depleted and thus more likely to experience hypotension following initiation of additional antihypertensive therapy. Whenever possible, all diuretics should be discontinued two to three days prior to the administration of telmisartan to reduce the likelihood

of hypotension. (See WARNINGS AND PRECAUTIONS, Hypotension). If this is not possible because of the patient's condition, telmisartan should be administered with caution and the blood pressure monitored closely. Thereafter, the dosage should be adjusted according to the individual response of the patient.

### **Patients with Renal Impairment**

The usual regimens of therapy with Auro-Telmisartan HCTZ may be followed as long as the patient's creatinine clearance is > 30 mL/min. In patients with more severe renal impairment, loop diuretics are preferred to thiazides; in this instance, Auro-Telmisartan HCTZ is not recommended.

### **Patients with Hepatic Impairment**

For patients with hepatic impairment, a starting dose of 40 mg of telmisartan is recommended. Auro-Telmisartan HCTZ is not recommended for patients with severe hepatic impairment.

### **Missed Dose**

If a dose is missed, patients should not take a double dose; patients should just carry on with the next dose at the usual time.

### **Administration**

Auro-Telmisartan HCTZ may be administered with or without food, however it should be taken consistently with regard to food intake.

## **OVERDOSAGE**

Limited information is available for telmisartan and hydrochlorothiazide tablets with regard to overdose in humans. Serum electrolytes and creatinine should be monitored frequently. If hypotension occurs, the patient should be placed in a supine position, with salt and volume replacements given quickly. For the individual components of telmisartan and hydrochlorothiazide tablets, the following information is available:

### ***Telmisartan***

Based on limited data, the most prominent manifestations of overdose are hypotension, dizziness and tachycardia; bradycardia also occurred in this setting as a result of parasympathetic (vagal) stimulation. If symptomatic hypotension should occur, supportive treatment should be instituted.

Telmisartan is not removed by hemodialysis.

### ***Hydrochlorothiazide***

The most common signs and symptoms observed are those caused by electrolyte depletion (hypokalemia, hypochloremia, hyponatremia) and dehydration resulting from excessive diuresis. If digitalis has also been administered, hypokalemia may accentuate cardiac arrhythmias.

The degree to which hydrochlorothiazide is removed by hemodialysis has not been established.

For management of a suspected drug overdose, contact your regional Poison Control Centre immediately.

## **ACTION AND CLINICAL PHARMACOLOGY**

### **Mechanism of Action**

Auro-Telmisartan HCTZ is a combination of telmisartan, a selective angiotensin II antagonist and hydrochlorothiazide, a thiazide diuretic.

#### ***Telmisartan***

Telmisartan is an orally active, AT<sub>1</sub> selective angiotensin II receptor antagonist. By selectively blocking the binding of angiotensin II to the AT<sub>1</sub> receptors, telmisartan inhibits the vasoconstrictor and aldosterone-secreting effects of angiotensin II. Telmisartan blocks AT<sub>1</sub> receptors, and has essentially no affinity for the AT<sub>2</sub> receptors. AT<sub>2</sub> receptors have been found in many tissues; to date, they have not been found to be associated with cardiovascular homeostasis.

Telmisartan does not inhibit angiotensin converting enzyme (ACE, also known as kininase II), the enzyme that converts angiotensin I to angiotensin II and degrades bradykinin, nor does it affect renin or other hormone receptors or ion channels involved in cardiovascular regulation of blood pressure and sodium homeostasis.

In hypertensive patients, antagonism of angiotensin II AT<sub>1</sub> receptors results in two to three-fold increases in plasma renin and angiotensin II plasma concentrations. Long term effects of increased AT<sub>2</sub> receptor stimulation by angiotensin II are unknown.

#### ***Hydrochlorothiazide***

Hydrochlorothiazide is a thiazide diuretic which affects the renal tubular mechanisms of electrolyte reabsorption, directly increasing excretion of sodium and chloride in the distal tubule, thus promoting water excretion. The diuretic action of hydrochlorothiazide reduces plasma volume, with consequent increases in plasma renin activity, increases in aldosterone secretion, increases in urinary potassium loss and decreases in serum potassium. The latter effects of the renin-aldosterone link are mediated by angiotensin II; as such, co-administration of an angiotensin II AT<sub>1</sub> receptor antagonist may prevent the potassium loss associated with thiazide diuretics. The precise mechanism of the antihypertensive effect of thiazides however, is not fully understood.

### **Pharmacodynamics**

#### ***Telmisartan***

The antihypertensive effects of telmisartan were demonstrated in 6 placebo-controlled clinical trials, in a total of 1773 patients, 1031 of whom were treated with telmisartan. Upon initiation of antihypertensive treatment with telmisartan, blood pressure was reduced after the first dose and there was a gradual increase in the antihypertensive effect during continued treatment for  $\leq 12$

weeks, with most of the increase occurring during the first month. Onset of antihypertensive activity occurs within 3 hours after administration of a single oral dose. The antihypertensive effect of once daily administration of telmisartan is maintained for the full 24- hour dose interval. The magnitude of blood pressure reduction from baseline, after placebo subtraction, was on average (SBP/DBP) -11.3/-7.3 mmHg for telmisartan 40 mg once daily, and -13.7/-8.1 mmHg for telmisartan 80 mg once daily. Upon abrupt cessation of treatment with telmisartan, blood pressure gradually returned to baseline values over a period of several days. During long term studies (without placebo control) the effect of telmisartan appeared to be maintained for  $\geq 1$  year.

For those patients treated with telmisartan 80 mg once daily who required additional blood pressure reduction, addition of a low dose of hydrochlorothiazide (12.5 mg) resulted in incremental blood pressure reductions of -9.4/-7.0 mmHg.

There was essentially no change in heart rate in telmisartan-treated patients in controlled trials.

In clinical trials with post-dose in- clinic monitoring no excessive blood pressure lowering peak effect was observed even after the first dose, and the incidence of symptomatic orthostasis was very low (0.04%).

With automated ambulatory blood pressure monitoring, the 24-hour trough-to-peak ratio for telmisartan was determined to be at least 80% for both systolic and diastolic blood pressure.

The antihypertensive effect of telmisartan is not influenced by patient age, weight or body mass index. Blood pressure in hypertensive black patients is significantly reduced by telmisartan (compared to placebo), but less so than in non-black patients.

In hypertensive patients with normal renal function, no clinically significant effects on renal plasma flow, filtration fraction, or glomerular filtration rate were observed. In multiple dose studies in hypertensive patients, telmisartan had no adverse effect on renal function as measured by serum creatinine or blood urea nitrogen.

*Diabetic Patients:* Multiple exploratory post hoc analyses were carried out on the three cardiovascular (CV) outcome trials (ONTARGET, TRANSCEND and PRoFESS). In TRANSCEND and PRoFESS, an increased risk of unexpected CV death was seen with telmisartan versus placebo in diabetics without previously diagnosed coronary artery disease (CAD) but not in those with a documented history of CAD. No such increased risk was demonstrated in ONTARGET for telmisartan versus ramipril in diabetes patients without previously diagnosed CAD.

These findings in diabetics with added cardiovascular risk, could be related to a pre-existing but asymptomatic or silent CAD. Diabetics with undiagnosed and therefore untreated CAD may be at increased risk when lowering blood pressure too far, e.g. when initiating antihypertensive therapy, due to a further reduction of perfusion in an already narrowed coronary artery.

### ***Hydrochlorothiazide***

After oral administration of hydrochlorothiazide, diuresis begins within 2 hours, peaks in

about 4 hours and lasts 6 to 12 hours.

***Telmisartan and Hydrochlorothiazide Combination***

In a placebo-controlled clinical study, the combination of telmisartan and hydrochlorothiazide resulted in decreases in trough systolic blood pressure (SBP) and diastolic blood pressure (DBP) which were greater than the decreases induced by either agent administered as monotherapy.

In a controlled clinical trial directly comparing telmisartan and hydrochlorothiazide tablets with telmisartan (80mg) monotherapy, trough SBP and DBP reductions observed with telmisartan and hydrochlorothiazide tablets were significantly greater than with telmisartan alone.

Similarly, in other controlled studies with patients who did not achieve or maintain adequate response with telmisartan monotherapy, the addition of 12.5 mg hydrochlorothiazide to titrated doses of telmisartan further reduced systolic and diastolic pressure.

The antihypertensive effect of telmisartan/hydrochlorothiazide (80 mg/12.5 mg) was independent of age or gender. The overall response to the combination was similar for black and non-black patients.

There was essentially no change in heart rate in patients treated with the combination of telmisartan and hydrochlorothiazide in the placebo-controlled trial.

**Pharmacokinetics**

There are no pharmacokinetic interactions between telmisartan and hydrochlorothiazide as the pharmacokinetic parameters of the individual components are unchanged by their co-administration as telmisartan and hydrochlorothiazide tablets. The results of a randomized, crossover study demonstrated that the bioavailabilities of telmisartan and hydrochlorothiazide were the same, whether administered as the fixed-dose combination or as the single entity formulations.

**Table 7: Single Dose Pharmacokinetics in Normotensive Subjects (10 Male and 10 Female Caucasian Subjects, 18 to 45 years of age). Given are arithmetic means (%CV).**

Drug	Therapy	C <sub>max</sub> (ng/mL)	t <sub>1/2</sub> (h)	AUC <sub>0-∞</sub> (ng·h/mL)	Clearance (CL/f) (mL/min.)	Volume of Distribution (V <sub>z</sub> /f) (L)
Telmisartan: monotherapy	A	246 (%CV 69.4)	22.2 (%CV 30)	1439 (%CV 94)	1650 (%CV 62)	2908 (%CV 60)
Telmisartan: combination therapy	B	266 (%CV 103)	24.4 (%CV 33)	1467 (%CV 94)	1565 (%CV 63)	3091 (%CV 63)
Hydrochlorothiazide: combination therapy	A	75.3 (%CV 26)	11.4 (%CV 43)	580.4 (%CV 27)	380 (%CV 23)	363.8 (%CV 43)
Hydrochlorothiazide: monotherapy	B	75.7 (%CV 22)	11.5 (%CV 36)	563.9 (%CV 20)	384 (%CV 20)	380.4 (%CV 40)

### ***Telmisartan***

**Absorption:** Following oral administration, telmisartan is well absorbed with a mean absolute bioavailability of about 50%. Mean peak plasma concentrations ( $C_{max}$ ) of telmisartan are reached in 0.5 - 1.0 hour after dosing. The pharmacokinetic profile is characterized by greater than proportional increases in plasma concentrations ( $C_{max}$  and AUC) with increasing doses >40 mg. Telmisartan shows bi-exponential decay kinetics with terminal elimination half life of approximately 24 hours, and does not accumulate in plasma upon repeated once daily administration. Food slightly reduces the bioavailability of telmisartan.

**Distribution:** Telmisartan is extensively bound to plasma proteins (>99.5%) at concentrations achieved at the recommended dosage. The apparent volume of distribution is approximately 500 L, suggesting extensive tissue binding sites.

**Metabolism:** Telmisartan is metabolized by conjugation to form a pharmacologically inactive acylglucuronide; this is the only metabolite that has been detected in human plasma and urine. Following both oral dosing and intravenous administration of radio labelled telmisartan, the parent compound represented approximately 85%, and the glucuronide approximately 11% of total radioactivity in plasma. The cytochrome P450 isoenzymes are not involved in the metabolism of telmisartan.

**Excretion:** Total plasma clearance of telmisartan is > 800 mL/min. Biliary excretion is the predominant route of elimination of telmisartan and its metabolite.

### ***Hydrochlorothiazide***

**Absorption:** Following oral administration, peak concentrations of hydrochlorothiazide were reached approximately 2.0 hours after dosing. Based on cumulative renal excretion of hydrochlorothiazide the absolute bioavailability was about 60% to 70%.

**Distribution:** Hydrochlorothiazide is 40% protein bound in the plasma and its apparent volume of distribution is 2 to 5 L/kg.

**Excretion:** Hydrochlorothiazide is not metabolized but is eliminated rapidly by the kidney. The plasma-half life has been observed to vary between 5.6 and 14.8 hours when the plasma levels can be followed for up to 24 hours. At least 61 percent of the oral dose is eliminated unchanged within 24 hours. Hydrochlorothiazide crosses the placenta but not the blood-brain barrier and is excreted in breast milk.

## **Special Populations and Conditions**

### ***Telmisartan***

#### **Pediatrics:**

Telmisartan pharmacokinetics have not been investigated in patients < 18 years of age.

#### **Geriatrics:**

The pharmacokinetics of telmisartan does not differ between elderly patients and those younger than 65 years of age.

**Gender:**

Plasma concentrations of telmisartan are generally 2-3 times higher in females than in males administered the same oral dose. In clinical trials however, no significant increases in blood pressure response or in the incidence of orthostatic hypotension were found in women. No dosage adjustment is necessary on the basis of gender.

**Hepatic Insufficiency:**

In patients with hepatic insufficiency, plasma concentrations of telmisartan are increased, and absolute bioavailability approaches 100% (see WARNINGS AND PRECAUTIONS and DOSAGE AND ADMINISTRATION). Reduction of the dose of telmisartan should be considered which would necessitate usage of the individual tablet formulations.

**Renal Insufficiency:**

Renal excretion of telmisartan is negligible. In patients with mild to moderate renal impairment, (creatinine clearance of 30-80 mL/min), no dosage adjustment is necessary (see WARNINGS AND PRECAUTIONS, Renal Impairment and DOSAGE AND ADMINISTRATION, Patients with Renal Impairment). Telmisartan is not removed by hemodialysis.

As a consequence of inhibiting the renin-angiotensin-aldosterone system, changes in renal function may be anticipated in susceptible individuals. In patients whose renal function may depend on the activity of the renin-angiotensin-aldosterone system such as patients with bilateral renal artery stenosis, unilateral renal artery stenosis to a solitary kidney or severe congestive heart failure, treatment with agents that inhibit this system has been associated with oliguria, progressive azotemia, and rarely, acute renal failure and/or death. In susceptible patients, concomitant diuretic use may further increase risk. Appropriate assessment of renal function should be conducted prior to use of Auro-Telmisartan HCTZ.

In studies of ACE inhibitors in patients with unilateral or bilateral renal artery stenosis, increases in serum creatinine or blood urea nitrogen were observed. Although there has been no long-term experience with telmisartan in this patient population, an effect similar to that observed with ACE inhibitors should be anticipated.

Due to the hydrochlorothiazide component, Auro-Telmisartan HCTZ is not recommended in patients with severe renal impairment (creatinine clearance  $\leq$  30 mL/min).

Thiazide diuretics should be used with caution in patients with renal impairment.

There is no experience regarding the administration of telmisartan and hydrochlorothiazide tablets in patients with a recent kidney transplant.

No initial dosing adjustment for telmisartan is necessary for elderly patients or for patients with renal impairment but greater sensitivity in some older individuals cannot be ruled out. Markedly reduced telmisartan plasma levels were observed in patients on hemodialysis.

The usual regimens of therapy with Auro-Telmisartan HCTZ may be followed as long as the patient's creatinine clearance is  $>30$  mL/min. In patients with more severe renal impairment,

loop diuretics are preferred to thiazides; in this instance Auro-Telmisartan HCTZ is not recommended.

Azotemia

Azotemia may be precipitated or increased by the hydrochlorothiazide component of Auro-Telmisartan HCTZ. Cumulative effects of the drug may develop in patients with impaired renal function since the primary route of excretion is through the urine.

If increasing azotemia and oliguria occur during treatment of severe progressive renal disease, the diuretic should be discontinued.

**Genetic Polymorphism:**

No studies were conducted to evaluate the influence of genetic polymorphisms on the pharmacokinetics or pharmacodynamics of telmisartan.

**STORAGE AND STABILITY**

Store at controlled room temperature (15°C to 30°C). Protect from moisture.

**DOSAGE FORMS, COMPOSITION AND PACKAGING**

Dosage form	Tablets	
	80 mg/ 12.5 mg	80 mg/25 mg
<b>Description</b>	White to off white and red coloured bi layered oblong shaped biconvex tablets debossed with “H” on one side and “72” on the other side. The white to off white layer may contain red specks.	White to off white and yellow coloured bi layered oblong shaped biconvex tablets debossed with “H” on one side and “76” on the other side. The white to off white layer may contain yellow specks.
<b>Composition</b>	Meglumine, povidone, sodium hydroxide pellets, colloidal silicon dioxide, mannitol, sodium stearyl fumarate, lactose monohydrate, hydroxypropyl cellulose & red ferric oxide.	Meglumine, povidone, sodium hydroxide pellets, colloidal silicon dioxide, mannitol, sodium stearyl fumarate, lactose monohydrate, hydroxypropyl cellulose & yellow ferric oxide.
<b>Packaging</b>	Blister pack: 3 x 10 Tablets HDPE Pack: 30’s, 100’s, 500’s & 1000’s Tablets	Blister pack: 3 x 10 Tablets HDPE Pack: 30’s, 100’s, 500’s & 1000’s Tablets



## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance - Telmisartan

*Proper name:* Telmisartan

*Chemical name:* 4'-[[4-Methyl-6-(1-methyl-1H-benzimidazol-2-yl)-2-propyl-1H-benzimidazol-1-yl]methyl]biphenyl-2-carboxylic acid


Or

4'-{[4-Methyl-6-(1-methyl-2-benzimidazolyl)-2-propyl-1-benzimidazolyl]methyl}-2-biphenyl carboxylic acid.

*Molecular formula:* C<sub>33</sub>H<sub>30</sub>N<sub>4</sub>O<sub>2</sub>,

*Molecular mass:* 514.6 g/mol

*Structural formula:*



*Physicochemical properties:*

**Description:**

A white or slightly yellowish, crystalline powder. Sparingly soluble in methylene chloride, slightly soluble in methanol, practically insoluble in water. It dissolves in 1M sodium Hydroxide.

**Polymorphism:**

Form A polymorphic form of telmisartan

Melting Point: 269 ± 1°C (polymorphic Form A)

log P = 7.73 ± 1.04

## Drug Substance - Hydrochlorothiazide


*Proper name:* Hydrochlorothiazide

*Chemical name:* 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide.

*Molecular formula:*  $C_7H_8ClN_3O_4S_2$

*Molecular mass:* 297.7 g/mol

*Structural formula:*



*Physicochemical properties:*

*Description:* Hydrochlorothiazide is a white or almost white, crystalline powder.  
In water - Very slightly soluble (1 part in 10000 parts)  
In acetone –Soluble (1 part in 30 parts)  
In alcohol - Sparingly soluble (1 part in 100 parts)  
In dilute solutions of alkali hydroxides - Dissolves

*Melting Point:* 274°C

## CLINICAL TRIALS

### Comparative bioavailability studies

#### **Telmisartan and Hydrochlorothiazide Tablets 80 mg/12.5 mg:**

A double blind, randomized, two-treatment, two-sequence, two-period, crossover, single-dose, comparative oral bioavailability study of 1 x 80 mg/12.5 mg Telmisartan and Hydrochlorothiazide Tablets USP (Test) of Aurobindo Pharma Limited, India manufactured for Auro Pharma Inc. (Canada) and 1 x 80/12.5 mg Micardis® Plus Tablets (Reference) of Boehringer Ingelheim (Canada) Ltd, Canada was conducted in 55 healthy, adult, male human, subjects under fasting conditions.

**Summary Table of Comparative Bioavailability Data**

<b>Telmisartan (1 x 80 mg Telmisartan and 12.5 mg Hydrochlorothiazide) From measured data Geometric Mean Arithmetic Mean (CV %)</b>				
<b>Parameter</b>	<b>Test*</b>	<b>Reference†</b>	<b>% Ratio of Geometric Means</b>	<b>90% Confidence Interval</b>
<b>AUC<sub>0→t</sub> (hr.ng/mL)</b>	2234.7 2665.4 (61.3)	2207.5 2670.5 (60.0)	101.2	97.0 – 105.7
<b>AUC<sub>0→∞</sub> (hr.ng/mL)</b>	2364.6 2842.5 (63.6)	2322.9 2816.7 (61.6)	101.8	97.7 – 106.1
<b>C<sub>max</sub> (ng/mL)</b>	439.0 495.5 (49.8)	462.9 536.7 (55.1)	94.8	84.3 – 106.7
<b>T<sub>max</sub>‡ (hr)</b>	1.0 (0.5-4.0)	1.0 (0.5-3.0)		
<b>T<sub>1/2</sub>§ (hr)</b>	16.5 (43.7)	16.2 (39.9)		

\*Auro-Telmisartan HCTZ (Telmisartan and Hydrochlorothiazide Tablets USP 80/12.5 mg), by Auro Pharma Inc.

†MICARDIS®PLUS Tablets 80/12.5 mg, of Boehringer Ingelheim (Canada) Ltd., Canada were purchased from Canada.

‡ Expressed as the median (range) only.

§ Expressed as arithmetic mean (%CV) only.

<b>Hydrochlorothiazide</b> <b>(1 x 80 mg Telmisartan and 12.5 mg Hydrochlorothiazide)</b> <b>From measured data</b> <b>Geometric Mean</b> <b>Arithmetic Mean (CV %)</b>				
<b>Parameter</b>	<b>Test*</b>	<b>Reference†</b>	<b>% Ratio of Geometric Means</b>	<b>90% Confidence Interval</b>
<b>AUC<sub>0→t</sub></b> <b>(hr.ng/mL)</b>	655.2 676.2 (25.2)	634.6 658.7 (26.7)	103.3	99.2-107.5
<b>AUC<sub>0→∞</sub></b> <b>(hr.ng/mL)</b>	734.6 757.0 (24.5)	712.6 738.7 (26.1)	103.1	99.2-107.2
<b>C<sub>max</sub> (ng/mL)</b>	97.8 100.5 (24.0)	103.3 107.6 (28.7)	94.7	90.5 – 99.0
<b>T<sub>max</sub>‡ (hr)</b>	1.8 (1.0-4.0)	1.8 (1.0-3.0)		
<b>T<sub>½</sub>§ (hr)</b>	8.4 (26.0)	8.1 (23.7)		

\*Auro-Telmisartan HCTZ (Telmisartan and Hydrochlorothiazide Tablets USP 80/12.5 mg), by Auro Pharma Inc.

†MICARDIS®PLUS Tablets 80/12.5 mg of Boehringer Ingelheim (Canada) Ltd., Canada were purchased from Canada.

‡ Expressed as the median (range) only.

§ Expressed as arithmetic mean (%CV) only.

### **Telmisartan and Hydrochlorothiazide Tablets 80 mg/25 mg:**

A double blind, randomized, two-treatment, two-sequence, two-period, crossover, single-dose, comparative oral bioavailability study of 1 x 80/25 mg Telmisartan and Hydrochlorothiazide Tablets USP (Test) of Aurobindo Pharma Limited, India manufactured for Auro Pharma Inc. (Canada) and 1 x 80/25 mg of Micardis® Plus Tablets (Reference) of Boehringer Ingelheim (Canada) Ltd, Canada was conducted in 51 healthy, adult, male human, subjects under fasting conditions.

### Summary Table of Comparative Bioavailability Data

<b>Telmisartan</b> <b>(1 x 80 mg Telmisartan and 25 mg Hydrochlorothiazide)</b> <b>From measured data</b> <b>Geometric Mean</b> <b>Arithmetic Mean (CV %)</b>				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval
<b>AUC<sub>0→t</sub></b> <b>(hr.ng/mL)</b>	2631.7 2953.6 (47.8)	2508.1 2894.3 (51.7)	104.9	99.8 - 110.3
<b>AUC<sub>0→∞</sub></b> <b>(hr.ng/mL)</b>	2781.4 3136.6 (48.9)	2660.2 3089.8 (53.9)	104.6	99.6 - 109.8
<b>C<sub>max</sub></b> (ng/mL)	532.6 593.8 (45.5)	491.1 571.0 (54.5)	108.5	97.1 - 121.1
<b>T<sub>max</sub></b> § (hr)	1.2 (0.3-3.0)	1.0(0.3-3.0)		
<b>T<sub>1/2</sub></b> § (hr)	17.9 (43.5)	17.3 (38.4)		

\*Auro-Telmisartan HCTZ (Telmisartan and Hydrochlorothiazide Tablets USP 80 /25 mg), by Auro Pharma Inc.

†MICARDIS®PLUS Tablets 80/25 mg, of Boehringer Ingelheim (Canada) Ltd., Canada were purchased from Canada.

§ Expressed as the median (range) only.

§ Expressed as arithmetic mean (%CV) only.

<b>Hydrochlorothiazide</b> <b>(1 x 80 mg Telmisartan and 25 mg Hydrochlorothiazide)</b> <b>From measured data</b> <b>Geometric Mean</b> <b>Arithmetic Mean (CV %)</b>				
Parameter	Test*	Reference†	% Ratio of Geometric Means	90% Confidence Interval
<b>AUC<sub>0→t</sub></b> <b>(hr.ng/mL)</b>	1444.2 1494.4 (27.2)	1513.1 1564.9 (25.7)	95.5	91.8 - 99.2
<b>AUC<sub>0→∞</sub></b> <b>(hr.ng/mL)</b>	1530.9 1577.9 (25.7)	1607.0 1654.1 (24.0)	95.3	91.9 - 98.7
<b>C<sub>max</sub></b> (ng/mL)	193.8 199.7 (24.4)	214.8 223.8 (28.0)	90.2	85.5 - 95.2
<b>T<sub>max</sub></b> § (hr)	1.8 (1.3-5.0)	1.5 (1.0-5.0)		
<b>T<sub>1/2</sub></b> § (hr)	9.0 (16.6)	9.2 (13.7)		

\*Auro-Telmisartan HCTZ (Telmisartan and Hydrochlorothiazide Tablets USP 80/25 mg), by Auro Pharma Inc.

†MICARDIS®PLUS Tablets 80/25 mg, of Boehringer Ingelheim (Canada) Ltd., Canada were purchased from Canada.

§ Expressed as the median (range) only.

§ Expressed as arithmetic mean (%CV) only.

## Study demographics and trial design

**Table 13- Summary of patient demographics for clinical trials**

Study #	Trial design	Dosage, route of administration and duration	Study subjects (n=number)	Mean age (Range)	Gender
502.261	Randomised, double-Blind	80 mg telmisartan and 80/12.5 mg telmisartan/hydrochlorothiazide, oral, 8 weeks	491 (245 telmisartan 80 mg; 246 telmisartan/hydrochlorothiazide (80/12.5 mg) patients)	55.3 (20 – 79)	males and females
502.204	Randomized, double-blind, placebo-controlled	Combinations of telmisartan & hydrochlorothiazide (T/H): 20/6.25, 20/12.5, 20/25, 40/6.25, 40/12.5, 40/25, 80/6.25, 80/12.5, 80/25, 160/6.25, 160/12.5/160/25 mg once daily, oral, 8 weeks	818	53.0* (19 – 80)	males and females
502.480	Randomized, double-blind, placebo-controlled trial in non-responders of the 80/12.5 mg strength	80 mg telmisartan/12.5 mg hydrochlorothiazide and 80 mg telmisartan/ 25 mg hydrochlorothiazide, oral, 8 weeks	713 (361 telmisartan/hydrochlorothiazide (80/12.5 mg); 352 telmisartan/hydrochlorothiazide (80/25 mg)	57.2 (28 – 93)	males and females

\* median age

T = telmisartan

H = hydrochlorothiazide

## Study results

In controlled clinical trials 571 patients were exposed to telmisartan 80 mg and concomitant hydrochlorothiazide 12.5 mg.

**Table 14**

Study #	Primary Measure of Efficacy	Patient Population	Summary of Results
502.261	Change from baseline in seated trough diastolic blood pressure (DBP) after 8 weeks of treatment.	Patients with mild to moderate hypertension who were taking no more than 3 antihypertensive agents at screening and who failed to respond adequately to telmisartan monotherapy	Treatment with telmisartan/hydrochlorothiazide fixed dose combination (80/12.5 mg) lowered trough DBP by an additional 3.1 mmHg and systolic blood pressure (SBP) by 5.7 mmHg compared to telmisartan 80 mg monotherapy. Reductions in both DBP and SBP were clinically and statistically significant (p<0.01).

Study #	Primary Measure of Efficacy	Patient Population	Summary of Results
502.204	Change from baseline in supine trough DBP after 8 weeks of treatment. Over 800 patients completed this study with approximately 70 patients in each of the primary dose groups	Patients with mild to moderate essential hypertension	The combination of telmisartan and hydrochlorothiazide (80/12.5 mg) was significantly ( $p<0.01$ ) better than either of its components administered as monotherapy in reducing trough supine diastolic blood pressure. Similar results were observed for supine systolic blood pressure and standing diastolic blood pressure.
502.480	Change from baseline in trough seated DBP after 8 weeks of treatment or at last trough observation during the double-blind treatment period	Patients without adequately controlled BP who failed to respond adequately to telmisartan/hydrochlorothiazide (80/12.5 mg)	Treatment with T80/H25 in patients with hypertension not adequately controlled by T80/H12.5 led to an additional, clinically relevant BP reduction. T80/H25 was superior to T80/H12.5 in reducing trough seated DBP after 8 weeks of randomised treatment. All analyses of secondary efficacy endpoints such as trough seated SBP, standing BP, and BP control and response showed better results for the T80/H25 group than for the T80/H12.5 group. Both treatments were safe and well tolerated.

BP = blood pressure

DBP = diastolic blood pressure

H = hydrochlorothiazide

SBP = systolic blood pressure

T = telmisartan

V = valsartan

**Table 15: Observed Mean Reduction from Baseline in Trough Supine Diastolic Blood Pressure (mmHg)**

	Placebo	Telmisartan 80 mg
Placebo	3.8	11.5 <sup>1</sup>
HCTZ 12.5 mg	7.3 <sup>1</sup>	14.9 <sup>2,3</sup>

<sup>1</sup> compared to placebo;  $p<0.01$

<sup>2</sup> compared to HCTZ 12.5 mg alone;  $p<0.01$

<sup>3</sup> compared to telmisartan 80 mg alone;  $p<0.01$

In 30 patients with mild to moderate hypertension, an 8-week randomized double-blind, parallel group comparison of the effect of telmisartan 80 mg versus telmisartan combined with hydrochlorothiazide (80/12.5 mg) on renal function in patients with mild to moderate hypertension was conducted. All patients randomized for the study had normal renal function. Renal parameters measured included: renal blood flow, effective renal plasma flow, glomerular filtration rate, urinary albumin, urinary protein, filtration fraction and renovascular resistance. Treatment with either regimen daily for 8 weeks did not significantly compromise renal function and there was no difference in renal function between telmisartan and telmisartan administered concomitantly with hydrochlorothiazide.

Study 502.480

At baseline, trough seated DBP means were comparable for both treatment groups with a mean of 95.0 mmHg for T80/H12.5 and 95.3 mmHg for T80/H25. Both groups showed a reduction in DBP by the end of study, with a larger reduction being observed in the T80/H25 treatment group. An adjusted mean change from baseline of -5.5 mmHg was observed for the T80/H12.5 group compared with an adjusted mean change of -7.1 mmHg for the T80/H25 group. The difference (95% CI) in the adjusted means of -1.6 mmHg (-2.5 mmHg,-0.6 mmHg) indicated the additional reduction in mean trough DBP present in the T80/H25 group. This difference was statistically significant (p=0.0012), which shows the superiority of T80/H25 over T80/H12.5 in reducing mean trough seated DBP in patients not responding adequately to T80/H12.5.

The analysis of the change from baseline in trough seated SBP at the last visit during the double-blind treatment phase was performed using the same methodology as for the primary endpoint. At baseline, trough seated SBP means of the 2 groups were comparable, with a mean of 147.4 mmHg for T80/H12.5 and 147.9 mmHg for T80/H25. Mean trough seated SBP decreased in both treatment groups by the end of study; the larger reduction was observed in the T80/H25 group. The adjusted mean change from baseline in the T80/H12.5 group was -7.1 mmHg, while for T80/H25 it was -9.8 mmHg. The difference (95% CI) between the groups in the adjusted means was -2.7 mmHg (-4.2 mmHg, -1.2 mmHg) with a p-value of 0.0003. The data show that treatment with T80/H25 reduced mean trough seated SBP more than T80/H12.5.

**Table 16: Analysis of Change from Baseline to the End of Study in Trough Seated DBP (FAS)**

Trough Seated DBP [mmHg]		T80/H12.5 N = 347	T80/H25 N = 340
Baseline:	Mean (SD)	95.0 (4.4)	95.3 (4.7)
End of Study:	Mean (SD)	89.6 (7.3)	88.3 (7.5)
	Adjust Mean* (SE)	89.6 (0.4)	88.0 (0.5)
Change to End of Study:	Mean (SD)	- 5.3 (6.4)	- 7.0 (6.8)
	Adjusted Mean* (SE)	- 5.5 (0.4)	- 7.1 (0.5)
Difference to T80/T12.5:	Adjusted Mean* (SE)	- 1.6 (0.5)	
	95% CI	(- 2.5, - 0.6)	
	p-value	0.0012	


\*Adjusted for baseline trough seated DBP and pooled country.

**Table 17: Analysis of Change from Baseline to the End of Study in Trough Seated SBP (FAS)**

Trough Seated SBP [mmHg]		T80/H12.5 N = 347	T80/H25 N = 340
Baseline:	Mean (SD)	147.4 (13.2)	147.9 (12.8)
End of Study:	Mean (SD)	141.8 (13.8)	139.5 (12.3)
	Adjust Mean* (SE)	140.6 (0.7)	137.9 (0.7)
Change to End of Study:	Mean (SD)	- 5.7 (11.0)	- 8.4 (10.6)
	Adjusted Mean* (SE)	- 7.1 (0.7)	- 9.8 (0.7)
Difference to T80/T12.5:	Adjusted Mean* (SE)	- 2.7 (0.7)	
	95% CI	(- 4.2, - 1.2)	
	p-value	0.0003	

\*Adjusted for baseline trough seated SBP and pooled country.





**Figure 1: Adjusted (for baseline and country) mean change from baseline (with SE) of trough seated BP in the pivotal trial 502.480**

In this study, both treatments were generally well tolerated as evidenced by a similar incidence of adverse event (AE) frequencies between T80/H12.5 (29.6%) and T80/H25 (31.5%) and the data obtained were consistent with the known safety profile of telmisartan and hydrochlorothiazide tablets. Frequently reported AEs in one or the other treatment arm (overall incidence  $\geq 1\%$  or 8 patients) were back pain (1.9% of the T80/H12.5 patients and vs. 2.0% of the T80/H25 patients), bronchitis (2.2% vs. 1.1%), headache (2.8% vs. 0.6%), palpitations (1.4% vs. 0.9%), and nasopharyngitis (0.6% vs. 1.7%). The frequency of AEs events considered drug related was also similar (5.0% for T80/H12.5 and 5.7% for T80/H25). Two serious AEs considered drug-related by the investigators were reported in the trial (atrioventricular block third degree in the T80/H12.5 group and atrial flutter in the T80/H25 group).

### Comparative Bioavailability Studies

An open-label, randomized, 4-way crossover, replicate-design study was conducted to compare the bioavailability following single oral administration of telmisartan (80 mg) and hydrochlorothiazide (12.5 mg) administered as either telmisartan and hydrochlorothiazide tablets or as the individual monotherapy formulations. Twenty (10M/10F) healthy subjects participated in this study. Comparative measures of exposure to telmisartan and hydrochlorothiazide were based on AUC and  $C_{max}$ . The results of this study demonstrated that both telmisartan and hydrochlorothiazide had similar bioavailabilities administered as either telmisartan and hydrochlorothiazide tablets or as the single entity formulations.

Conventional average bioequivalence, average scaled bioequivalence, and individual bioequivalence (as a secondary analysis) using a moment-based scaled approach were evaluated with respect to Telmisartan pharmacokinetic variables. With respect to HCTZ pharmacokinetic variables, conventional average bioequivalence was evaluated. The upper bounds of the one-sided 95% confidence interval for the average, scaled bioequivalence measure  $M_{as}$  are, on the ratio scale, 109.4% and 113.8% for the primary pharmacokinetic variables  $AUC_{0-\infty}$  and  $C_{max}$ , respectively. These upper bounds of the confidence intervals are below the upper bound of the bioequivalence range, 125%. Thus average, scaled bioequivalence with respect to the variables  $AUC_{0-\infty}$  and  $C_{max}$  is shown. Similarly, average, scaled bioequivalence with respect to the secondary variable  $AUC_{0-48h}$  was shown. The 90% confidence intervals for the “test/reference” mean ratio for the primary pharmacokinetic variables  $AUC_{0-\infty}$  and  $C_{max}$  are 100.0% to 111.0%

and 106.8% to 129.0%, respectively. The confidence interval for  $AUC_{0-\infty}$  falls in the bioequivalence range of 80% to 125%, while the confidence interval for  $C_{max}$  falls in the bioequivalence range of 75% to 133%. The confidence interval for the “test/reference” mean ratio of the secondary variable  $AUC_{0-48h}$  falls in the 80% to 125%, bioequivalence range.

**Table 18: Results for Telmisartan**

<p><b>Telmisartan (80 mg)</b></p> <p><b>From measured data</b></p> <p><b>Adjusted Geometric Mean</b></p> <p><b>Arithmetic Mean (CV %)</b></p>
---

<b>Parameter</b>	<b>Fixed dose combination (Test)*</b>	<b>Individual tablets (Reference)<sup>†</sup></b>	<b>% Ratio of Adjusted Geometric Means</b>	<b>90% Confidence Interval</b>
$AUC_{0-48h}$ (ng·h/ml)	875.9 1158 (85.51)	820.7 1082 (83.46)	106.7	101.1 - 112.7
$AUC_{0-\infty}$ (ng·h/ml)	1034.5 1414 (92.50)	981.9 1364 (91.97)	105.4	100.0 - 111.0
$C_{MAX}$ (ng/mL)	196.4 251.5 (87.88)	167.3 207.0 (71.58)	117.4	106.8 - 129.0
$T_{MAX}^{\S}$ (h)	1.00 (0.50 - 2.00)	1.00 (0.50 - 2.00)		
$T_{1/2}^{  }$ (h)	24.12 (29.65)	24.73 (35.90)		

\*batch no. 9960326

<sup>†</sup>Telmisartan<sup>®</sup> 80 mg oblong tablet, batch no. 9960326;

<sup>§</sup>Expressed as the median (range) only.

<sup>||</sup>Expressed as the arithmetic mean (CV%) only.

For Hydrochlorothiazide, the 90% confidence intervals for the “test/reference” mean ratio for the primary pharmacokinetic variables  $AUC_{0-\infty}$  and  $C_{max}$  are 91.8% to 109.5% and 90.8% to 107.4%, respectively. The confidence intervals for the “test/reference” mean ratio of the pharmacokinetic variable  $Ae_{0-48h}$  is 91.0% to 103.9%. All confidence intervals fall in the bioequivalence range of 80% to 125%.

**Table 19: Results for Hydrochlorothiazide**

<b>Hydrochlorothiazide (12.5 mg) From measured data</b>  <b>Adjusted Geometric Mean Arithmetic Mean (CV %)</b>
--

Parameter	Fixed dose combination (Test)*	Individual tablets (Reference)†	% Ratio of Adjusted Geometric Means	90% Confidence Interval
AUC <sub>0-24h</sub> (ng·h/ml)	478.1 493.9 (25.73)	467.2 475.04 (19.09)	102.3	97.3 - 107.6
AUC <sub>0-∞</sub> (ng·h/ml)	575.4 580.4 (27.46)	574.0 563.9 (19.85)	100.3	91.8 - 109.5
C <sub>MAX</sub> (ng/mL)	73.1 75.33 (26.14)	74.1 75.72 (26.61)	98.7	90.8 - 107.4
T <sub>MAX</sub> § (h)	2.00 (1.00 -3.00)	2.00 (1.00 -3.00)		
T <sub>1/2</sub>    (h)	11.35 (43.10)	11.51 (36.33)		

\*batch no. 9960326

†Hydrochlorothiazide 12.5 mg tablet, batch no. .F4260;

§Expressed as the median (range) only.

||Expressed as the arithmetic mean (CV%) only.

## DETAILED PHARMACOLOGY

### Animal

The effect of telmisartan in combination with hydrochlorothiazide was investigated in spontaneously hypertensive rats. Repeated oral administration of telmisartan at a dose of 3 mg/kg/day for 5 days to conscious rats reduced mean arterial blood pressure (MAP) significantly and persistently with maximal decrease in MAP of approximately 36 mmHg.

Hydrochlorothiazide alone (10 mg/kg/day) had no effect on blood pressure in this model, however when administered in combination with telmisartan (3 mg/kg/day), induced a significantly greater antihypertensive effect than with telmisartan alone, with a maximal reduction of about 53 mmHg. Furthermore, the telmisartan/hydrochlorothiazide combination ameliorated the alteration in potassium balance when compared to hydrochlorothiazide alone in this model.

A slight, significant increase in heart rate (~ 20 bpm) was observed during treatment with telmisartan and hydrochlorothiazide in combination; this increase reverted to control values during the washout period.

## Human

### **Drug Interaction - Telmisartan and Hydrochlorothiazide**

A randomized, 3-way crossover study was conducted in 14 healthy subjects to investigate the potential for a pharmacokinetic interaction between telmisartan and hydrochlorothiazide when administered concomitantly. Oral doses of either hydrochlorothiazide alone (25 mg, qd x 7), telmisartan alone (160 mg, qd x 7) or both drugs at the respective doses in combination, daily for 7 days were administered. Plasma concentrations of both telmisartan and hydrochlorothiazide were assessed at steady state. Based on a comparative analysis, it was concluded that there is no pharmacokinetic interaction between telmisartan and hydrochlorothiazide when administered concomitantly.

### **Bioavailability**

See CLINICAL TRIALS, Comparative Bioavailability Studies.

## **TOXICOLOGY**

### **Chronic Toxicity**

#### ***Telmisartan and Hydrochlorothiazide***

Repeated-dose toxicity studies of 26 weeks duration were conducted in both rats and dogs. These studies were designed to compare the toxicological profiles of telmisartan and hydrochlorothiazide administered alone, with that of the drugs given in combination.

**Table 20: Repeat Dose Toxicity Studies Conducted with Telmisartan/Hydrochlorothiazide**

<b>Species</b>	<b>Dose Telmisartan/HCTZ (mg/kg/day)</b>	<b>Duration</b>	<b>NOTEL (mg/kg/day)</b>
Rat (Chbb:THOM) (20/sex/dose)	0/0 0.1/0.03 4/1.2 50/7.8 50/15.6 50/0 0/15.6	26 weeks	0.1/0.03
Dogs (Beagle) (4/sex/dose)	0/0 0.25/0.08 1/0.31 4/0.63 4/1.25 4/0 0/1.25	26 weeks	0.25/0.08

HCTZ = hydrochlorothiazide

Repeated, oral doses of telmisartan with and without hydrochlorothiazide for 26 weeks in rats induced a pronounced and persistent dose-related decrease of blood pressure without reflex tachycardia. At 50 mg/kg of telmisartan, the addition of hydrochlorothiazide had an additive effect on the blood pressure-lowering effect of telmisartan. Clinical laboratory and histopathological changes were similar to those observed in previous toxicity studies in rats with

telmisartan alone. Essentially, there were no new toxicities observed with the addition of hydrochlorothiazide.

In dogs, repeated oral doses of telmisartan with hydrochlorothiazide administered for 26 weeks was associated with nephrotoxicity, which is consistent with the findings from previous studies with telmisartan alone. The addition of 0.63 mg/kg of hydrochlorothiazide to 4 mg/kg telmisartan did not increase renal toxicity whereas the co-administration of 1.25 mg/kg hydrochlorothiazide significantly increased toxicity. The exacerbation of nephrotoxicity in this species can be ameliorated by saline supplementation.

### **Reproductive Toxicity**

***Telmisartan and hydrochlorothiazide:*** A developmental toxicity study was conducted in rats with oral doses of telmisartan and hydrochlorothiazide used in combination (3.2/1.0, 15/4.7, 50/15.6 and 0/15.6 mg/kg/day). Although the two higher dose combinations appeared to be more toxic to the dams than either drug alone, the results indicated the lack of teratogenic, fetotoxic or embryotoxic potential of the telmisartan/hydrochlorothiazide combination at the doses tested in this animal model.

***Telmisartan:*** In studies on fertility and reproductive performance in male and female rats, no effect on mating performance, reproductive organs, or fertility in either sex or on litter parameters was observed with telmisartan oral doses of 5-100 mg/kg. No teratogenic or embryotoxic potential in rats was observed at oral doses of  $\leq 50$  mg/kg administered during gestation. However, at toxic dose levels, non-clinical studies indicated some hazardous potential of telmisartan to fetal development (increased number of late resorptions in rabbits) and to the postnatal development of the offspring: lower body weight, delayed eye opening, and higher mortality.

Telmisartan was detectable in the placenta, fetus and amniotic fluid of rats after single oral doses of 1 mg/kg.

***Hydrochlorothiazide:*** Hydrochlorothiazide was orally administered to pregnant mice and rats during gestation, at doses of up to 3000 and 1000 mg/kg/day respectively. There was no evidence of fetotoxicity or teratogenicity.

Hydrochlorothiazide had no adverse effects on the fertility of mice and rats of either sex in studies where these species were exposed via their diet, to doses of  $\leq 100$  and  $\leq 4$  mg/kg respectively, prior to mating and throughout gestation.

### **Carcinogenicity and Mutagenicity**

No carcinogenicity or mutagenicity studies have been conducted with the combination of telmisartan and hydrochlorothiazide. However, these studies have been conducted for telmisartan and hydrochlorothiazide alone. Based on the preclinical safety profile of the telmisartan and hydrochlorothiazide combination and on human pharmacokinetic studies, there is no indication of any adverse interaction between telmisartan and hydrochlorothiazide.

### ***Telmisartan***

The carcinogenic potential of telmisartan was assessed in 2-year feeding studies in mice at doses of 10, 100 and 1000 mg/kg/day in rats at 3, 15, and 100 mg/kg/day. Drug administration did not affect overall survival time in either study nor did it affect the rate of tumour-induced mortality. There were no increases in overall tumour incidence, incidence of benign or malignant tumours or in tumour multiplicity associated with telmisartan administration.

The standard battery of genotoxicity assays did not reveal any telmisartan -related effects at either the gene or chromosome level. These assays included bacterial mutagenicity tests with *Salmonella typhimurium* and *E. coli*, a gene mutation test with CHO cells, a cytogenetic test with human lymphocytes and an *in vivo* mouse micronucleus assay.

### ***Hydrochlorothiazide***

The carcinogenic potential of hydrochlorothiazide was assessed in 2-year feeding studies in mice at doses of up to 600 mg/kg/day and rats at doses of up to 100 mg/kg/day. There was no evidence of carcinogenic potential in female mice or male and female rats; equivocal evidence for hepatocarcinogenicity in male mice was reported.

Hydrochlorothiazide was not genotoxic in the Ames mutagenicity assay with *Salmonella typhimurium* strains TA 98, TA 100, TA 1535, TA 1537 and TA 1538, in the Chinese Hamster Ovary (CHO) test for chromosomal aberrations or *in vivo* assays using mouse germ cell chromosomes, Chinese hamster bone marrow chromosomes and the *Drosophila* sex-linked recessive lethal trait gene. Positive test results were obtained only in the *in vitro* CHO Sister Chromatid Exchange (clastogenicity) and in the Mouse Lymphoma Cell (mutagenicity) assays using concentrations of hydrochlorothiazide from 43 to 1300 mcg/mL and in the *Aspergillus nidulans* non-disjunction assay at unspecified concentrations.

## REFERENCES

McGill JB, Reilly PA. Combination treatment with telmisartan and hydrochlorothiazide in black patients with mild to moderate hypertension. *Clin Cardiol* 2001; 24(1):66-72.

Neutel JM, Smith DHG. Dose response and antihypertensive efficacy of the AT1 receptor antagonist telmisartan in patients with mild to moderate hypertension. *Adv Ther* 1998;15(4):206-217.

Young CL, Dias VC, Stangier J. Multiple-dose pharmacokinetics of telmisartan and of hydrochlorothiazide following concurrent administration in healthy subjects. *J Clin Pharmacol* 2000; 40 (12 Pt 1):1323-30.

Product monograph – MICARDIS® PLUS, Boehringer Ingelheim (Canada) Ltd, Date of Revision: February 17, 2017, Submission Control No: 200158.

**IMPORTANT: PLEASE READ**

**PART III: CONSUMER INFORMATION**

**PrAuro-Telmisartan HCTZ**

**Telmisartan and Hydrochlorothiazide Tablets, USP**

80 mg/12.5 mg and 80 mg/25 mg

**Read this carefully before you start taking Auro-Telmisartan HCTZ and each time you get a refill. This leaflet is a summary and will not tell you everything about Auro-Telmisartan HCTZ. Talk to your doctor, nurse, or pharmacist about your medical condition and treatment and ask if there is any new information about Auro-Telmisartan HCTZ.**

**ABOUT THIS MEDICATION**

**What the medication is used for:**

Auro-Telmisartan HCTZ is a combination of 2 drugs. It is used when 2 drugs are required to treat your high blood pressure.

**What it does:**

Auro-Telmisartan HCTZ contains a combination of 2 drugs, telmisartan and hydrochlorothiazide:

- Telmisartan is an angiotensin receptor blocker (ARB). You can recognize an ARB because its medicinal ingredient ends in “-SARTAN”. It lowers blood pressure.
- Hydrochlorothiazide is a diuretic or “water pill” that increases urination. This lowers blood pressure.

This medicine does not cure high blood pressure. It helps to control it. Therefore, it is important to continue taking Auro-Telmisartan HCTZ regularly even if you feel fine.

**When it should not be used:**

Do not take Auro-Telmisartan HCTZ if you:

- Are allergic to telmisartan, hydrochlorothiazide or to any non-medical ingredient in the formulation.
- Are allergic to any sulfonamide-derived drugs (sulfa drugs), most of them have a medicinal ingredient that ends in “-MIDE”. This includes other diuretics (“water pills”).
- Have experienced an allergic reaction (angioedema) with swelling of the hands, feet or ankles, face, lips, tongue, throat, or sudden difficulty breathing or swallowing to any ARB (any drug in the same class as Telmisartan). Be sure to tell your doctor, nurse, or pharmacist that this has happened to you.
- Have difficulty urinating or produce no urine.
- Are pregnant or intend to become pregnant. Taking Auro-Telmisartan HCTZ during pregnancy can cause

injury and even death to your baby.

- Are breastfeeding. Auro-Telmisartan HCTZ passes into breast milk.
- Are allergic to some sugars (fructose, lactose and/or mannitol).
- Are already taking a blood pressure-lowering medicine that contains aliskiren (such as Rasilez) and you have diabetes or kidney disease.
- Have one of the following rare hereditary diseases:
  - o Galactose intolerance;
  - o Lapp lactase deficiency;
  - o Glucose-galactose malabsorption.

Because lactose is a non-medicinal ingredient in Auro-Telmisartan HCTZ.

**What the medicinal ingredients are:**

Telmisartan and hydrochlorothiazide

**What the non-medicinal ingredients are:**

Meglumine, povidone, sodium hydroxide pellets, colloidal silicon dioxide, mannitol, sodium stearyl fumarate, lactose monohydrate, hydroxypropyl cellulose, yellow ferric oxide (80 mg/25 mg) & red ferric oxide (80 mg/12.5 mg).

**What dosage forms it comes in:**

Tablets 80 mg/12.5 mg and 80 mg/25 mg.

**WARNINGS AND PRECAUTIONS**

**Serious Warnings and Precautions –Pregnancy Auro-Telmisartan HCTZ should not be used during pregnancy. If you discover that you are pregnant while taking Auro-Telmisartan HCTZ, stop the medication and contact your doctor, nurse or pharmacist as soon as possible.**

Before you use Auro-Telmisartan HCTZ, talk to your doctor, nurse or pharmacist if you:

- Are allergic to any drug used to lower blood pressure including angiotensin converting enzyme (ACE) inhibitors, or penicillin.
- Have narrowing of an artery or a heart valve.
- Have heart failure.
- Have diabetes, liver or kidney disease.
- Are taking a medicine that contains aliskiren, such as Rasilez, used to lower high blood pressure. The combination with Auro-Telmisartan HCTZ is not recommended.
- Are taking an angiotensin-converting-enzyme inhibitor (ACEI).
- Have lupus or gout.
- Are on dialysis.
- Are dehydrated or if you suffer from excessive vomiting, diarrhea, or sweating.



## IMPORTANT: PLEASE READ

- Are taking a salt substitute that contains potassium, potassium supplements, or a potassium-sparing diuretic (a specific kind of “water pill”).
- Are on a low-salt diet.
- Are less than 18 years old.
- Are having surgery and general anesthesia, (even at the dentist's office), tell the doctor or dentist that you are taking Auro-Telmisartan HCTZ as there may be a sudden fall in blood pressure associated with general anesthesia.
- Have been told by your doctor that you have an intolerance to some sugars.

### Hydrochlorothiazide in Auro-Telmisartan HCTZ can cause Sudden Eye Disorders:

- Myopia: sudden nearsightedness or blurred vision.
- Glaucoma: an increased pressure in your eyes, eye pain.

Untreated, it may lead to permanent vision loss. These eye disorders are related and can develop within hours to weeks of starting Auro-Telmisartan HCTZ.

You may become sensitive to the sun while taking Auro-Telmisartan HCTZ. Exposure to sunlight should be minimized until you know how you respond.

**Driving and using machines:** Before you perform tasks which may require special attention, wait until you know how you respond to Auro-Telmisartan HCTZ. Dizziness, light headedness, or fainting can especially occur after the first dose and when the dose is increased.

### INTERACTIONS WITH THIS MEDICATION

As with most medicines, interactions with other drugs are possible. Tell your doctor, nurse, or pharmacist about all the medicines you take, including drugs prescribed by other doctors, vitamins, minerals, natural supplements, or alternative medicines.

The following may interact with Auro-Telmisartan HCTZ:

- Adrenocorticotrophic hormone (ACTH) used to treat West Syndrome.
- Alcohol, barbiturates (sleeping pills), or narcotics (strong pain medications). They may cause low blood pressure and dizziness when you go from lying or sitting to standing up.
- Amphotericin B, an antifungal drug.
- Anticancer drugs, including cyclophosphamide and methotrexate.
- Antidepressants, in particular selective serotonin reuptake inhibitors (SSRIs), including citalopram, escitalopram, and sertraline.
- Antidiabetic drugs, including insulin and oral medicines.
- Bile acid resins used to lower cholesterol, such as Cholestyramine and Colestipol Resins.
- Other blood pressure lowering drugs, including diuretics (“water pills”), aliskiren-containing products (e.g. Rasilez), or angiotension-converting-enzyme inhibitors (ACEI). When taken in combination with Auro-Telmisartan HCTZ, they may cause excessively low blood pressure.
- Calcium or vitamin D supplements.
- Corticosteroids used to treat joint pain and swelling.
- Digoxin, a heart medication.
- Drugs that slow down or speed up bowel function, including atropine, metoclopramide, and domperidone.
- Drugs used to treat epilepsy, including carbamazepine and topiramate.
- Gout medications, including allopurinol and probenecid.
- Lithium used to treat bipolar disease.
- Nonsteroidal anti-inflammatory drugs (NSAIDs), used to reduce pain and swelling. Examples include ibuprofen, naproxen, and celecoxib.
- Skeletal muscle relaxants used to relieve muscle spasms, including tubocurare.
- Pressor Amines (e.g. norepinephrine)
- Warfarin.

### PROPER USE OF THIS MEDICATION

Take Auro-Telmisartan HCTZ exactly as prescribed. It is recommended to take your dose at about the same time everyday preferably in the morning.

Auro-Telmisartan HCTZ can be taken with or without food, but it should be taken the same way each day. If Auro-Telmisartan HCTZ causes upset stomach, take it with food or milk.

#### Usual adult dose:

The recommended dose of Auro-Telmisartan HCTZ is one tablet daily.

#### Overdose:

If you think you have taken too much Auro-Telmisartan HCTZ contact your healthcare professional, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

#### Missed dose:

If you have forgotten to take your dose during the day, carry on with the next one at the usual time. Do not

**IMPORTANT: PLEASE READ**

double dose.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Side effects may include:

- back or leg pain, muscle cramps, joint pain, muscle spasms, pain, weakness, restlessness
- headache, anxiety, dizziness, pins and needles in your fingers
- diarrhea, constipation, nausea, vomiting, upset stomach, abdominal pain, flatulence, decreased appetite, enlargement of the glands in your mouth
- dry mouth
- rash, eczema, skin eruptions, bleeding under the skin, red patches on the skin
- drowsiness, insomnia, fatigue
- visual disturbances
- upper respiratory infection
- reduced libido
- reduction in blood platelets, which increases risk of bleeding or bruising (small purple-red marks in skin or other tissue caused by bleeding)
- low blood magnesium level
- high blood calcium level
- increased PH (disturbed acid-base balance) due to low blood chloride level

**If any of these affects you severely, tell your doctor, nurse or pharmacist.**

**Auro-Telmisartan HCTZ can cause abnormal blood test results. Your doctor will decide when to perform blood tests and will interpret the results.**

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
<b>Common</b>	<b>Low Blood Pressure:</b> dizziness, fainting, light headedness May occur when you go from lying or sitting to standing up.	√		
	<b>Anemia:</b> fatigue, loss of energy, weakness, shortness of breath		√	

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>				
Symptom / effect		Talk to your healthcare professional		Stop taking drug and get immediate medical help
		Only if severe	In all cases	
<b>Uncommon</b>	<b>Electrolyte Imbalance:</b> weakness, drowsiness, muscle pain or cramps, irregular heartbeat		√	
	<b>Chest pain</b>		√	
	<b>Kidney Disorder:</b> change in frequency of urination, nausea, vomiting, swelling of extremities, fatigue		√	
	<b>Decreased White Blood Cells:</b> infections, fatigue, fever, aches, pains, and flu-like symptoms		√	
	<b>Increased blood sugar:</b> frequent urination, thirst, and hunger	√		
<b>Rare</b>	<b>Urinary tract infection (Cystitis):</b> frequent or painful urination, feeling unwell		√	
	<b>Depression:</b> Low mood, loss of interest in activities, change in appetite and sleep patterns	√		
	<b>Decreased or increased levels of potassium in the blood:</b> Irregular heartbeats, muscle weakness and generally feeling unwell		√	

**IMPORTANT: PLEASE READ**

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
<b>Liver disorder:</b> Yellowing of the skin or eyes, dark urine, abdominal pain, nausea, vomiting, loss of appetite		√	
<b>Eye disorders:</b> - <b>Myopia:</b> sudden near sightedness or blurred vision - <b>Glaucoma:</b> increased pressure in your eyes, eye pain			√
<b>Low blood sugar:</b> shaky, irregular heartbeat, sweating, hunger, dizziness		√	
<b>Decreased Platelets:</b> bruising, bleeding, fatigue and weakness		√	
<b>Unknown</b>			√
<b>Allergic Reaction:</b> Rash, hives, swelling of the face, lips, tongue or throat, difficulty swallowing or breathing			√
<b>Rhabdomyolysis:</b> Muscle pain that you cannot explain, muscle tenderness or weakness, dark brown urine			√
<b>Heart Rhythm or Heart Rate disturbances:</b> heart racing or skipping a beat		√	
<b>Sepsis (blood poisoning):</b> chills, confusion, fever or low body		√	

<b>SERIOUS SIDE EFFECTS, HOW OFTEN THEY HAPPEN AND WHAT TO DO ABOUT THEM</b>			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
temperature, shakiness, irregular heartbeat (including fatal outcome)			
<b>Toxic Epidermal Necrolysis:</b> severe skin peeling, especially in mouth and eyes			√
<b>Inflammation of the Pancreas:</b> abdominal pain that lasts and gets worse when you lie down, nausea, vomiting		√	

*This is not a complete list of side effects. For any unexpected effects while taking Auro-Telmisartan HCTZ, contact your doctor or pharmacist.*

**HOW TO STORE IT**

Store at controlled room temperature (15°C to 30°C).  
Protect from moisture.

**Store Auro-Telmisartan HCTZ out of the reach and sight of children and pets.**

**IMPORTANT: PLEASE READ**

**Reporting Side Effects**

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<http://www.hc-sc.gc.ca/dhpmpps/medeff/report-declaration/index-eng.php>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

*NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.*

**MORE INFORMATION**

**If you want more information about Auro-Telmisartan HCTZ:**

- Talk to your healthcare professional
- Find the full product monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<http://hc-sc.gc.ca/index-eng.php>); the manufacturer's website [www.auropharma.ca](http://www.auropharma.ca), or by calling 1-855-648-6681.

This leaflet was prepared by

**Auro Pharma Inc.**  
3700 Steeles Avenue West, Suite # 402  
Woodbridge, Ontario, L4L 8K8,  
Canada.

Date of Revision: April 18, 2017